

The modular variety of hyperelliptic curves of genus three

Eberhard Freitag
Mathematisches Institut
Im Neuenheimer Feld 288
D69120 Heidelberg
 FREITAG@
 MATHI.UNI-HEIDELBERG.DE

Riccardo Salvati Manni
Dipartimento di Matematica
Piazzale Aldo Moro, 2
I-00185 Roma, Italy
 SALVATI@
 MAT.UNIROMA1.IT

2007

Introduction

The modular variety of non singular and complete hyperelliptic curves with level-two structure of genus 3 is a 5-dimensional quasi projective variety which admits several standard compactifications. The first one comes from the period map, which realizes this variety as a sub-variety of the Siegel modular variety of level two and genus three $\mathcal{H}_3/\Gamma_3[2]$. We denote the hyperelliptic locus by $\mathcal{I}_3[2]$ and its closure in the Satake compactification by

$$\overline{\mathcal{I}_3[2]} \subset \overline{\mathcal{H}_3/\Gamma_3[2]}.$$

The hyperelliptic locus has 36 irreducible (isomorphic) components, which correspond to the 36 even theta characteristics. We denote the component of the characteristic m by

$$X = \overline{\mathcal{I}_3[2]}^m.$$

One of the purposes of this paper will be to describe the equations of X in a suitable projective embedding and its Hilbert function (5.8). It will turn out that X is normal.

Two further models use the fact that hyperelliptic curves of genus three can be obtained as coverings of a projective line with 8 branch points. The level two structure of the curve corresponds to an ordering of these 8 points. This leads to the configuration space of 8 ordered points on a line. There are two other important compactifications of this configuration space. The first one comes from geometric invariant theory using so-called semistable degenerated point configurations in $(P^1)^8$. We denote this GIT-compactification by Y . The equations of this variety in a suitable projective embedding (by so-called Specht polynomials) are known. For a general approach to the GIT-model $(P_1)^n // \mathrm{SL}(2)$ we refer to the paper [HM1], [HM2]. In the special case $n = 8$ this variety also can be identified with a Baily-Borel compactified ball-quotient [Ko2].

$$Y = \overline{\mathcal{B}/\Gamma[1-i]}.$$

We will describe these results in some detail and obtain new proofs including some finer results for them. We will see for example that the graded algebra describing $(P_1)^8 // \mathrm{SL}(2)$ is a Gorenstein ring (1.4) and that it describes the full ring of modular forms of $\Gamma[1 - i]$ (2.4). This part is essentially due to Kondo [Ko2], but we can avoid the theory of K_3 -surfaces which is an essential part of Kondo's approach.

The other compactification uses the fact that families of marked projective lines $(P^1, x_1, \dots, x_8)$ can degenerate to stable marked curves $(C, x_1, \dots, x_8)$ of genus 0.

We use the standard notation $\bar{M}_{0,8}$ for this compactification. It is known that $\bar{M}_{0,n}$ for arbitrary n is a smooth and projective fine moduli scheme.

In [Ka] (see also [AL]) it has been proved that the birational map

$$\bar{M}_{0,n} \longrightarrow (P_1)^n // \mathrm{SL}(2)$$

is everywhere regular. It is also known that there is natural everywhere regular map (Torelli map)

$$\bar{M}_{0,n} \longrightarrow \overline{\mathcal{H}_g / \Gamma_g[2]} \quad (\text{Satake compactification})$$

with $n = 2g + 2$. Hence in the case $n = 8$ we have a diagram

$$\begin{array}{ccc} & \bar{M}_{0,8} & \\ \swarrow & & \searrow \\ Y & \text{---} & X \end{array}$$

The horizontal arrow is only birational but not everywhere regular.

In this paper we find another realization of this triangle which uses the fact that there are graded algebras (closely related to algebras of modular forms) A, B such that

$$X = \mathrm{proj}(A), \quad Y = \mathrm{proj}(B).$$

The rational map $Y \text{ --- } X$ is induced by a homomorphism of graded algebras

$$A \longrightarrow B.$$

This homomorphism rests on the theory of Thomae (19th century), in which the thetanullwerte of hyperelliptic curves have been computed. We use Mumfords approach [Mu] for the construction of this homomorphism. Using the explicit equations for A, B we can compute the base locus of $Y \rightarrow X$. It turns out to be the union of 56 P^3 (6.2). Blowing up the base locus we get a projective variety Y^* and a diagram

$$\begin{array}{ccc} & Y^* & \\ \swarrow & & \searrow \\ Y & \text{---} & X \end{array},$$

where the diagonal arrows are everywhere regular. But the model Y^* is singular. The space Y contains 35 special points (corresponding to non-stable points in the GIT-model or to the cusps in the ball-quotient model). Their inverse images in Y^* are 35 disjoint rational surfaces. Blowing up them we get a dominant *smooth* model $\tilde{Y} \rightarrow Y^*$ and especially a diagram

$$\begin{array}{ccc} & \tilde{Y} & \\ \swarrow & & \searrow \\ Y & \text{-----} & X. \end{array}$$

We will see that $\tilde{Y}$ and $\bar{M}_{0,8}$ are isomorphic, 6.7. This can be considered as an explicit description of $\bar{M}_{0,8}$.

The variety $\tilde{Y}$ gives a correspondence between subvarieties of $Y = \bar{X}(8)^{\text{GIT}}$ and of $Y = \bar{\mathcal{B}}/\Gamma[1-i]$. This correspondence explains several combinatorial similarities between the two models. These similarities can be described best, if one uses the ball-model to describe Y . The reason is that in the description of the Siegel and the ball-model the space $\mathbb{F}_2^6$ occurs. In the Siegel case it occurs as a symplectic space, its elements are so-called characteristics. In the ball case it occurs as quadratic space. The symmetry of both models can be explained by the fact that in characteristic 2 the symplectic and orthogonal world come together. This will be described in the last section.

The second named author is grateful to Corrado De Concini and Shigeyuki Kondo for helpful and enlightening discussions. He would like also to thank Herbert Lange and Hanspeter Kraft for bringing his attention to the papers [AL] and [Ho].

1. The configuration space for eight points on a line

We recall some basic facts about the configuration space of eight points on the line. With the help of computer computations we get very easy proofs of them and also some new insight.

We consider the subset

$$\mathcal{X}(8) \subset P^1(\mathbb{C})^8$$

of ordered eight tuples of elements of the projective line, which are pairwise distinct. The elements of $P^1(\mathbb{C})$ can be represented by columns $\begin{pmatrix} a \\ b \end{pmatrix}$, which are different from zero and which are defined up to a constant factor. Hence the elements of $\mathcal{X}(8)$ can be represented by matrices

$$M := \begin{pmatrix} a_1 & a_2 & \dots & a_8 \\ b_1 & b_2 & \dots & b_8 \end{pmatrix}.$$

Since $\mathrm{SL}(2, \mathbb{C})$ acts on $P^1(\mathbb{C})$, we get an action of $\mathrm{SL}(2, \mathbb{C})$ on $\mathcal{X}(8)$ by multiplication from the left. The configuration space is defined as

$$X(8) = \mathrm{SL}(2, \mathbb{C}) \backslash \mathcal{X}(8).$$

We recall that a tableau is a matrix

$$\begin{pmatrix} i_1 & \dots & i_4 \\ j_1 & \dots & j_4 \end{pmatrix}$$

which contains as entries all digits i , $1 \leq i \leq 8$, and with the property

$$i_1 < i_2 < i_3 < i_4, \quad i_1 < j_1, \quad i_2 < j_2, \quad i_3 < j_3, \quad i_4 < j_4.$$

There are 105 tableaux. The tableau is called standard if in addition $j_1 < j_2 < j_3 < j_4$. There are 14 standard tableaux:

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 & 5 \\ 4 & 6 & 7 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 & 6 \\ 4 & 5 & 7 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 3 & 7 \\ 4 & 5 & 6 & 8 \end{pmatrix} \\ \begin{pmatrix} 1 & 2 & 4 & 5 \\ 3 & 6 & 7 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 4 & 6 \\ 3 & 5 & 7 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 4 & 7 \\ 3 & 5 & 6 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 5 & 6 \\ 3 & 4 & 7 & 8 \end{pmatrix} \\ \begin{pmatrix} 1 & 2 & 5 & 7 \\ 3 & 4 & 6 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 3 & 4 & 5 \\ 2 & 6 & 7 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 3 & 4 & 6 \\ 2 & 5 & 7 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 3 & 4 & 7 \\ 2 & 5 & 6 & 8 \end{pmatrix} \\ \begin{pmatrix} 1 & 3 & 5 & 6 \\ 2 & 4 & 7 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 & 8 \end{pmatrix}$$

To each tableau one associates the expression

$$D(M) = \prod_{\nu=1}^4 (a_{i_\nu} b_{j_\nu} - a_{j_\nu} b_{i_\nu}).$$

This leads to the so-called Specht polynomial

$$D \begin{pmatrix} 1 & \dots & 1 \\ X_1 & \dots & X_8 \end{pmatrix} = (X_{i_1} - X_{j_1})(X_{i_2} - X_{j_2})(X_{i_3} - X_{j_3})(X_{i_4} - X_{j_4}).$$

The space generated by all the 105 Specht polynomials has dimension 14. A basis is given by the Specht polynomials of the standard tableaux. We denote them by $Y_1, \dots, Y_{14}$ in the above ordering. It is clear that these polynomials define a holomorphic map

$$X(8) \longrightarrow P^{13}(\mathbb{C}).$$

It is easy to check and well-known that $X(8) \hookrightarrow P^{13}$ is a smooth embedding. We denote by $\bar{X}(8)$ the closure of the image of $X(8)$ in $P^{13}(\mathbb{C})$. Hence

$$\bar{X}(8) = \mathrm{proj} \, \mathbb{C}[Y_1, \dots, Y_{14}] / \mathcal{J},$$

where $\mathcal{J}$ is the ideal of relations between the Specht polynomials. The ideal $\mathcal{J}$ can be computed by elimination: Consider the polynomial ring in 28 variables $\mathbb{C}[X_1, \dots, X_{14}, T_1, \dots, T_{14}]$ and the ideal generated by the relations $T_i - Y_i$ ($1 \leq i \leq 14$). Then $\mathcal{J}$ is the intersection of this ideal with $\mathbb{C}[Y_1, \dots, Y_{14}]$. With computer algebra one can verify now*) a slightly modified result of Koike [Koi]:

*) The computer algebra calculation has to be done over the field $\mathbb{Q}$ instead $\mathbb{C}$, which is sufficient.

1.1 Lemma (Koike). *The ideal $\mathcal{J}$ is generated by the polynomials*

$$\begin{aligned}
J_1 &= Y_9Y_{13} - Y_8Y_{14} \\
J_2 &= Y_2Y_{12} - Y_3Y_{12} - Y_5Y_{12} + Y_6Y_{12} - Y_1Y_{14} + Y_3Y_{14} + Y_5Y_{14} - Y_7Y_{14} - \\
&\quad Y_8Y_{14} + Y_9Y_{14} - Y_{11}Y_{14} + Y_{13}Y_{14} \\
J_3 &= Y_7Y_{11} - Y_6Y_{12} \\
J_4 &= Y_4Y_{11} - Y_3Y_{12} - Y_4Y_{13} + Y_3Y_{14} \\
J_5 &= Y_2Y_{11} - Y_5Y_{11} - Y_3Y_{12} + Y_6Y_{12} - Y_1Y_{13} + Y_5Y_{13} - Y_7Y_{13} + Y_3Y_{14} - \\
&\quad Y_{11}Y_{14} + Y_{13}Y_{14} \\
J_6 &= Y_7Y_{10} - Y_9Y_{10} - Y_5Y_{12} + Y_5Y_{14} \\
J_7 &= Y_6Y_{10} - Y_8Y_{10} - Y_5Y_{11} + Y_8Y_{11} - Y_9Y_{11} + Y_5Y_{13} - Y_6Y_{13} + Y_6Y_{14} \\
J_8 &= Y_4Y_{10} - Y_9Y_{10} - Y_3Y_{12} - Y_5Y_{12} + Y_6Y_{12} - Y_4Y_{13} - Y_1Y_{14} + \\
&\quad Y_3Y_{14} + Y_4Y_{14} + Y_5Y_{14} - Y_7Y_{14} + Y_{10}Y_{14} - Y_{11}Y_{14} + Y_{14}^2 \\
J_9 &= Y_3Y_{10} - Y_8Y_{10} - Y_5Y_{11} + Y_8Y_{11} - Y_9Y_{11} - Y_3Y_{12} + Y_6Y_{12} - Y_1Y_{13} + \\
&\quad Y_5Y_{13} - Y_7Y_{13} + Y_{10}Y_{13} - Y_{11}Y_{13} + Y_3Y_{14} + Y_{13}Y_{14} \\
J_{10} &= Y_2Y_7 - Y_3Y_7 - Y_1Y_9 + Y_3Y_9 - Y_9Y_{11} - Y_5Y_{12} + Y_6Y_{12} + Y_5Y_{14} - \\
&\quad Y_7Y_{14} + Y_9Y_{14} \\
J_{11} &= Y_4Y_6 - Y_3Y_7 - Y_4Y_8 + Y_7Y_8 + Y_3Y_9 - Y_6Y_9 - Y_7Y_{13} + Y_6Y_{14} \\
J_{12} &= Y_2Y_6 - Y_3Y_7 - Y_1Y_8 + Y_3Y_9 - Y_5Y_{11} - Y_9Y_{11} + Y_6Y_{12} + Y_5Y_{13} - \\
&\quad Y_7Y_{13} + Y_8Y_{14} \\
J_{13} &= Y_4Y_5 - Y_3Y_7 - Y_4Y_8 + Y_7Y_8 - Y_1Y_9 + Y_3Y_9 + Y_4Y_9 - Y_7Y_9 - Y_9Y_{11} - \\
&\quad Y_5Y_{12} + Y_6Y_{12} - Y_7Y_{13} + Y_5Y_{14} + Y_9Y_{14} \\
J_{14} &= Y_3Y_5 - Y_3Y_7 - Y_1Y_8 + Y_3Y_9 - Y_5Y_{11} - Y_9Y_{11} + Y_6Y_{12} + Y_5Y_{13} - \\
&\quad Y_7Y_{13} + Y_8Y_{14}
\end{aligned}$$

We want to describe the boundary $\bar{X}(8) - X(8)$. Koike compares $\bar{X}(8)$ with a certain ball-quotient. This relies to a paper of Kondo who identifies the Satake compactified ball-quotient with the GIT-compactification of $X(8)$. Since Kondo's argument is not completely correct (s. section 2), we give here in some detail a self contained complete description of $\bar{X}(8)$. We need some results from geometric invariant theory which gives the so-called GIT-compactification of $X(8)$.

Recall that this is a certain projective variety $\text{proj } R^G$. Here R is the graded algebra $R = \sum_{n=0}^{\infty} R_n$, where R_n denotes the space of all polynomials in the entries of a 2×8 -matrix M which are homogenous of the same degree n in each column of M . Hence $\text{proj}(R) = P^1(\mathbb{C})^8$. There is a natural action of the group $G = \text{SL}(2, \mathbb{C})$ on R . The ring of invariants R^G is finitely generated. One defines

$$\bar{X}(8)^{\text{GIT}} := \text{proj}(R^G).$$

Recall that a point from $P^1(\mathbb{C})^8$ is called semistable, if there is some element from R_n^G , which doesn't vanish on it. The set of all semistable points $P^1(\mathbb{C})_{ss}^8 \subset P^1(\mathbb{C})^8$ is a G -invariant Zariski open subset. There is an obvious map $P^1(\mathbb{C})_{ss}^8 \rightarrow \bar{X}(8)^{\text{GIT}}$ and this is a categorical quotient. Since we are in a geometric situation, this map is surjective.

It is known that a point in $P^1(\mathbb{C})^8$ is semistable if and only if not more than 4 of its components can agree. From this description it follows that the Specht polynomials don't have a joint zero. Hence we have a map $P^1(\mathbb{C})_{ss}^8 \rightarrow \bar{X}(8)$ and from the universal property we get a map

$$\bar{X}(8)^{\text{GIT}} \longrightarrow \bar{X}(8).$$

We will see that this is a biholomorphic map. This actually follows from a known theorem from Kempe, s. [HM1], [HM2]:

The ring R^G is generated by its elements of degree one. (These are the Specht polynomials).

We don't need to use this theorem. We will obtain it as consequence of a sharper result which seems to be new (1.4).

First of all one sees from the description of the semistable points that the irreducible subvarieties of dimension 4 of the boundary $\bar{X}(8)^{\text{GIT}} - X(8)$ are defined by the subsets of $P^1(\mathbb{C})^8$ where two points agree. Hence these irreducible subsets perform one orbit under the group S_8 . It follows that the boundary $\bar{X}(8) - X(8)$ also contains at most one orbit of irreducible subvarieties of dimension 4. It is really one orbit, since one can check easily by means of computer algebra that the subvariety defined by $J_9 = J_{10} = J_{11} = J_{12} = J_{13} = J_{14} = 0$ is of dimension 4. It is easy to exhibit a special point in this subvariety which is smooth in the whole $\bar{X}(8)$. Hence we see:

1.2 Lemma. *The ring $\mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}$ is regular in codimension one.*

Another calculation with computer algebra gives an explicit minimal resolution of this ring:

1.3 Lemma. *Let A be the polynomial ring $\mathbb{C}[Y_1, \dots, Y_{14}]$. The minimal resolution of the ring $B := \mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}$ is of the form*

$$\begin{aligned} 0 \rightarrow A^1 \rightarrow A^{14} \rightarrow A^{175} \rightarrow A^{512} \rightarrow A^{700} \\ \rightarrow A^{512} \rightarrow A^{175} \rightarrow A^{14} \rightarrow A^1 \rightarrow B \rightarrow 0. \end{aligned}$$

Corollary. *The ring B is a Cohen Macaulay ring, even more, it is a Gorenstein ring.*

Using Serre's criterion for normality we get now:

1.4 Theorem. *The ring $\mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}$ is a normal Gorenstein ring.*

Corollary 1 (Kempe). *This ring agrees with the full ring of invariants R^G which hence is generated by the elements of lowest degree (hence by the Specht polynomials).*

Corollary 2 (Kempe). *The map*

$$\bar{X}(8)^{\text{GlT}} \xrightarrow{\sim} \bar{X}(8)$$

is biholomorphic.

We want to point out that Kempe's result has been generalized in [HM1] and [HM2] to point configurations of an arbitrary number of points in the projective line.

From the computation of the resolution one can also read off:

1.5 Theorem. *The Hilbert function of $R^G \cong \mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}$ is*

$$\begin{aligned} \sum_{n=0}^{\infty} \dim R_n^G t^n &= \frac{1 + 8t + 22t^2 + 8t^3 + t^4}{(1-t)^6} \\ &= 1 + 14t + 91t^2 + 364t^3 + 1085t^4 + 5719t^6 + \dots \end{aligned}$$

This formula is in concordance with other formulae in literature. Howe [Ho] for example gives

$$\dim R_n^G = \frac{n^5 + 5n^4 + 11n^3 + 13n^2 + 9n + 3}{3},$$

which is obviously the same. The fact that Howe's formula is true for all n , reflects that R^G is Gorenstein.

Another remarkable formula has been communicated to us by De Concini

$$\begin{aligned} \dim R_n^G &= \sum_{i_1+2i_2+\dots+ni_n=4n} \binom{8}{i_1} \binom{8-i_1}{i_2} \binom{8-i_1-\dots-i_n}{i_n} \\ &\quad - \sum_{i_1+2i_2+\dots+ni_n=4n-1} \binom{8}{i_1} \binom{8-i_1}{i_2} \binom{8-i_1-\dots-i_n}{i_n} \end{aligned}$$

2. A ball quotient

The ring $\mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}$ of the previous section turns out to be the ring of modular forms on a certain ball quotient. This is related to the fact that $\bar{X}(8)$ is isomorphic to a Baily-Borel compactified ball-quotient. These results are essentially due to Kondo [Ko2]. Here we amend and extend some of his arguments.

The configuration space of 8 points on a line is related to the even lattice

$$L = U \oplus U(2) \oplus D_4(-1) \oplus D_4(-1).$$

Here as usual $U = \mathbb{Z}^2$ with the quadratic form x_1x_2 and

$$D_4 = \{ (x_1, x_2, x_3, x_4) \in \mathbb{Z}^4; \quad x_1 + \dots + x_4 \equiv 0 \pmod{2} \}, \quad q(x) = x_1^2 + \dots + x_4^2.$$

The notation $M(m)$ means that one takes the same abelian group M , but the quadratic form is multiplied by m . Hence $U \oplus U(2) = \mathbb{Z}^4$ with the quadratic form $x_1x_2 + 2x_3x_4$. The discriminant group L'/L is isomorphic to $\mathbb{F}_2^6$. The isomorphism can be chosen such that the diagram

$$\begin{array}{ccc} L'/L & \xrightarrow{q} & \mathbb{Q}/\mathbb{Z} \\ \uparrow & & \uparrow \\ \mathbb{F}_2^6 & \longrightarrow & \mathbb{F}_2 \\ (x_1, \dots, x_6) & \longmapsto & x_1x_4 + x_2x_5 + x_3x_6 \end{array}$$

commutes. Here $\mathbb{F}_2$ is embedded into $\mathbb{Q}/\mathbb{Z}$ by sending $0 \mapsto 0$ and $1 \mapsto 1/2$. It will be basic for us that the orthogonal group (so-called even type) $O(\mathbb{F}_2^6)$ is isomorphic to the symmetric group S_8 . We choose an isomorphism

$$S_8 \xrightarrow{\sim} O(\mathbb{F}_2^6).$$

A natural concrete construction of such an isomorphism will be given in section 4. At the moment the choice is not important.

We now describe a certain graded algebra, which is related to $O(\mathbb{F}_2^6)$. This is the third member of a sequence of algebras related to $O(\mathbb{F}_2^{2m})$ [FS]. We recall very briefly its definition. We attach to each maximal totally isotropic subspace $I \subset \mathbb{F}_2^{2m}$ (of dimension m) a variable X_I and we consider the subring

$$\mathcal{R}_m := \mathbb{C}[\dots X_I - X_J \dots]$$

of the polynomial ring $\mathbb{C}[\dots X_I \dots]$ in all these variables. We define an ideal $\mathcal{I}_m \subset \mathcal{R}_m$. It is the sum of a linear part and a quadratic part,

$$\mathcal{I}_m = \mathcal{I}_m^{\text{lin}} + \mathcal{I}_m^{\text{qua}}.$$

We first define the linear part: For this we need the characteristic function

$$\chi_I : \mathbb{F}_2^{2m} \longrightarrow \mathbb{C}$$

of a maximal totally isotropic subspace. Let $A \subset \mathbb{F}_2^{2m}$ be a totally isotropic subspace of dimension $m - 2$. There are 6 maximal totally isotropic subspaces $I_1, \dots, I_6$ containing A . Their ordering can be chosen such that

$$(\chi_{I_1} - \chi_{I_2}) + (\chi_{I_3} - \chi_{I_4}) + (\chi_{I_5} - \chi_{I_6}) = 0.$$

The linear part $\mathcal{I}_m^{\text{lin}}$ then is generated by all

$$(X_{I_1} - X_{I_2}) + (X_{I_3} - X_{I_4}) + (X_{I_5} - X_{I_6}).$$

Next we define the quadratic part: Again we consider a totally isotropic subspace $A \subset \mathbb{F}_2^{2m}$ of dimension $m - 2$ and the 6 maximal totally isotropic subspaces $I_1, \dots, I_6$ containing it. After a suitable choice of their ordering the relation

$$(\chi_{I_1} - \chi_{I_2})(\chi_{I_1} - \chi_{I_4}) = (\chi_{I_3} - \chi_{I_6})(\chi_{I_5} - \chi_{I_6})$$

holds. The quadratic part $\mathcal{I}_m^{\text{qua}}$ of the ideal is generated by all

$$(X_{I_1} - X_{I_2})(X_{I_1} - X_{I_4}) - (X_{I_3} - X_{I_6})(X_{I_5} - X_{I_6}).$$

The ring $\mathcal{R}_m/\mathcal{I}_m$ seems to be very interesting. For example the cases $m = 5$ and $m = 6$ are related to Enriques surfaces. Here we are interested in the case $m = 3$.

First of all one computes that a linear combination $\sum_V C_V X_V$ is in the ideal $\mathcal{I}_3$ if and only if $\sum_V C_V \chi_V = 0$. The dimension of the space generated by all $\chi_V - \chi_W$ can be computed and it is 14. Hence $\mathcal{R}_3/\mathcal{I}_3^{\text{lin}}$ is a polynomial ring in 14 variables. The group $O(\mathbb{F}_2^6)$ acts on this ring. The action comes from an irreducible linear representation on the lowest degree part. Since the subgroup of index two $A_8 \subset S_8$ admits only one irreducible representation of dimension 14, we obtain an essentially unique A_8 -equivariant isomorphism

$$\mathcal{R}_3/\mathcal{I}_3^{\text{lin}} \longrightarrow \mathbb{C}[Y_1, \dots, Y_{14}],$$

where on the right hand side we have the ring of section one. It turns out to be $O(\mathbb{F}_2^6) = S_8$ -equivariant. Now a concrete check shows that the quadratic relations go to 0. Hence we get a homomorphism

$$\mathcal{R}_3/\mathcal{I}_3 \longrightarrow \mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}.$$

It is no problem to check by a concrete calculation:

2.1 Theorem. *The homomorphism*

$$\mathcal{R}_3/\mathcal{I}_3 \longrightarrow \mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}$$

is an $O(\mathbb{F}_2^6) \cong S_8$ -equivariant isomorphism.

Corollary. *The ring $\mathcal{R}_3/\mathcal{I}_3$ is a normal ring of Krull dimension 5.*

We want to mention that the dimension of $\mathcal{R}_m/\mathcal{I}_m$ is unknown for $m > 3$.

The ring $\mathcal{R}_3/\mathcal{I}_3$ is related to modular forms [FS]: There is a subgroup of index two of $O(L \otimes_{\mathbb{Z}} \mathbb{R})$, which doesn't contain the reflections along vectors of positive norm. Its intersection with $O(L)$ is denoted by $O'(L)$. The discriminant kernel is

$$\Gamma_L = \text{kernel}(O'(L) \longrightarrow O(L'/L)).$$

The Borchers additive lifting attaches to each maximal totally isotropic subspace a modular form [FS]. This gives a homomorphism

$$\mathcal{R}_3/\mathcal{I}_3^{\text{lin}} \longrightarrow A(\Gamma_L),$$

where $A(\Gamma_L)$ denotes the ring of modular forms with respect to Γ_L . It is possible to derive certain quadratic relations [FS]. Kondo derived quartic relations but Koike [Koi] pointed out that they are consequences of quadratic relations. These relations can be explained naturally by means of the theory of Borchers products: To explain this, we have to recall the notion of a star [FS].

2.2 Definition. *A star S is a set of 4 anisotropic vectors in $\mathbb{F}_2^6$ which is a coset of a 2-dimensional totally isotropic space.*

(Kondo considers instead of stars “maximal totally singular subspaces”. They are in one-to one- correspondence with the stars. The set of anisotropic elements of such a space is a star and this gives a one-to-one correspondence between stars and maximal totally singular subspaces.) To every star a certain Borchers product can be associated. To explain this we have to recall how to associate to an element $\alpha \in \mathbb{F}_2^6$ of non-zero norm a certain Heegner divisor $\mathcal{H}_\alpha$ on the symmetric domain which is associated to $O(L)$. Recall that the elements of this symmetric domain can be considered as two dimensional positive definite subspaces of $L \otimes_{\mathbb{Z}} \mathbb{R}$. Then $\mathcal{H}_\alpha$ consists of all such subspaces, which are orthogonal to some

$$\delta \in L^*, \quad (\delta, \delta) = -1, \quad \delta \text{ over } \alpha.$$

If S is star, we denote by $\mathcal{H}_S$ the union of the four $\mathcal{H}_\alpha$, $\alpha \in S$. The basic fact is that the additive lift space contains for every star S a form f_S , whose zero divisor contains $\mathcal{H}_S$. Since there is also a Borchers product of the same weight with this property, the zero divisor equals $\mathcal{H}_S$. From this it is possible to derive directly quadratic relations [FS] and to prove:

2.3 Lemma. *The Borchers additive lifting gives a homomorphism*

$$\mathcal{R}_3/\mathcal{I}_3 \longrightarrow A(\Gamma_L).$$

Of course this homomorphism is not an isomorphism, since the left hand side has Krull dimension 6 and the right hand side has Krull dimension 11. The picture remedies if one intersects Γ_L with a certain unitary group:

The lattice L admits a complex structure. We follow Kondo's description [Ko2]. For this one uses the isomorphism $\mathbb{Z}^4 \rightarrow \mathbb{Z}^4$ defined by the matrix

$$\begin{pmatrix} -1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 2 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & -1 \end{pmatrix}.$$

This is an isometry

$$\rho_1 : U \oplus U(2) \xrightarrow{\sim} U \oplus U(2)$$

We also use the isometry

$$\rho_0 : D_4 \xrightarrow{\sim} D_4, \quad (x_1, x_2, x_3, x_4) \longmapsto (x_2, -x_1, x_4, -x_3).$$

The direct sum $\rho := \rho_1 \oplus \rho_0 \oplus \rho_0$ defines an isometry

$$\rho : L \xrightarrow{\sim} L.$$

Since $\rho^2 = -\text{id}$ we get a structure as $\mathbb{Z}[i]$ module on L by defining

$$ia := \rho(a).$$

One has $L \cong \mathbb{Z}[i]^6$. We denote by $(a, b) = q(a+b) - q(a) - q(b)$ the real bilinear form on L . Because of $(a, \rho(a)) = 0$ the pairing

$$2\langle a, b \rangle := (a, b) - i(a, ib) \quad (\implies 2\langle a, a \rangle = (a, a)).$$

is a hermitian form of signature $(1, 5)$. Notice that we take hermitian forms to be $\mathbb{C}$ -linear in the second variable. Let $U(L, \langle \cdot, \cdot \rangle)$ denote the unitary group. We set

$$\Gamma := O'(L) \cap U(L, \langle \cdot, \cdot \rangle), \quad \Gamma[1-i] := \Gamma_L \cap U(L, \langle \cdot, \cdot \rangle).$$

The natural homomorphism

$$\Gamma/\Gamma[1-i] \longrightarrow O'(L)/\Gamma_L \longrightarrow O(\mathbb{F}_2^6) \cong S_8$$

is still an isomorphism.

We denote by $\mathcal{B}$ the associated complex five dimensional ball. It can be considered as the set of positive definit one-dimensional complex subspaces of $L \otimes_{\mathbb{Z}} \mathbb{R}$. We can intersect the Heegner divisor $\mathcal{H}_\alpha$ with the ball:

$$\mathcal{B}_\alpha := \mathcal{H}_\alpha \cap \mathcal{B}.$$

The group Γ acts on $\mathcal{B}$ and one can consider the graded algebras of modular forms $A(\Gamma[1-i])$. There is a natural restriction homomorphism $A(\Gamma_L) \rightarrow A(\Gamma[1-i])$. Together with 2.3 we get a homomorphism $\mathcal{R}_3/\mathcal{I}_3 \rightarrow A(\Gamma[1-i])$.

2.4 Theorem. *The homomorphism*

$$\mathcal{R}_3/\mathcal{I}_3 \rightarrow A(\Gamma[1-i])$$

is an isomorphism. Hence this ring of modular forms is generated by 14 modular forms with 14 defining quadratic relations. The dimension formula is given by 1.5.

Corollary. *There is an induced biholomorphic map*

$$\bar{X}(8) \longrightarrow \overline{\mathcal{B}/\Gamma[1-i]},$$

where the right hand side denotes the Baily-Borel compactification.

This result can be found in Kondo's paper [Ko2]. But the proof is not convincing. Kondo constructs by means of the theory of $K3$ -surfaces a period map $X(8) \rightarrow \mathcal{B}/\Gamma[1-i]$. Then he wants to use an extension theorem of Borel to extend it to the compactifications. This extension theorem of Borel states essentially that a holomorphic map of a punctured disk into an arithmetic quotient D/Γ extends to a holomorphic map of the full disk to the Satake compactification. The theorem of Borel requires that Γ is torsion free. Otherwise it is false as for example $\mathrm{SL}(2, \mathbb{Z})$ shows. Also $\Gamma[1-i]$ can be taken as counter example.

The group $\Gamma[1-i]$ is not torsion free. If r is an element with $2\langle r, r \rangle = (r, r) = -2$ then the complex reflection

$$x \longmapsto x + 2\langle x, r \rangle x = x + (x, r)x - i(x, ir)x$$

is contained in $\Gamma[1-i]$. These reflections are related to the divisors $\mathcal{B}_\alpha = \mathcal{H}_\alpha \cap \mathcal{B}$. By definition this divisor consists of all $z \in \mathcal{B}$ such that

$$(\delta, z) = (\delta, iz) = 0 \quad \text{for all } \delta \in L^*, \quad (\delta, \delta) = -1, \quad \text{over } \alpha.$$

Using the hermitian form this is equivalent to $\langle \delta, z \rangle = 0$. This we can rewrite as

$$\langle r, z \rangle = 0, \quad r := (1-i)\delta.$$

Obviously $r \in L$ and $\langle r, r \rangle = -1$. This shows that $\mathcal{B}_\alpha$ is the fixed point set of a reflection inside $\Gamma[1-i]$.

2.5 Remark. *The zero divisor of the restriction $F_S = f_S|_{\mathcal{B}}$ of the star modular form is contained in the ramification divisor of the projection $\mathcal{B} \rightarrow \mathcal{B}/\Gamma[1-i]$.*

Because of the ramification the star modular forms F_S , restricted to the ball, vanish at $\mathcal{B}_\alpha$, $\alpha \in S$ of order two. The ramification makes another correction in [Ko2] necessary. It is possible to take a holomorphic square root G_S of F_S . These forms are modular forms with characters. Hence for two stars S_1, S_2 the quotient G_{S_1}/G_{S_2} is not invariant under $\Gamma[1-i]$. The stars and the tableaux are in 1-1 correspondence [Ko2]. Denote by τ_1, τ_2 the tableaux which correspond to S_1, S_2 and by $\mu_{\tau_1}, \mu_{\tau_2}$ the corresponding Specht polynomials. Kondo claims $G_{S_1}/G_{S_2} = \mu_{\tau_1}/\mu_{\tau_2}$ (s. the two lines before theorem 7.6 in [Ko2]). The correct equation is

$$F_{S_1}/F_{S_2} = \mu_{\tau_1}/\mu_{\tau_2}.$$

The point is that the pole- and zero orders of F_{S_1}/F_{S_2} considered as functions on the *quotient* $\mathcal{B}/\Gamma[1-i]$ are only one and not two.

Proof of theorem 2.4. From the description of the zeros of the F_S it follows that they have no common zero on the Baily-Borel compactified ball quotient. Hence one obtains a finite map

$$\overline{\mathcal{B}/\Gamma[1-i]} \longrightarrow \text{proj}(\mathcal{R}_3/\mathcal{I}_3).$$

Since $\mathcal{R}_3/\mathcal{I}_3$ is normal, one has only to show that this map is generically injective. This follows from Kondo's comparison from $X(8)$ and $\mathcal{B}/\Gamma[1-i]$ by means of K3-surfaces. It can also be proven directly using the methods of [FS]. In this way one can avoid the use of K3-surfaces. (It depends on the situation and also on a question of taste whether one wants to apply the theory of moduli to the theory of modular forms or conversely.)

3. Siegel modular forms of genus three

Let

$$\mathcal{H}_g = \{ Z \in \mathbb{C}^{(g,g)}; \quad Z = {}^t Z, \text{ Im } Z > 0 \}$$

be the Siegel half plane of genus g and

$$\Gamma_g[q] = \text{kernel}(\text{Sp}(g, \mathbb{Z}) \longrightarrow \text{Sp}(g, \mathbb{Z}/q\mathbb{Z}))$$

the principal congruence subgroup of level q in the Siegel modular group $\Gamma_g := \Gamma_g[1]$. It acts on $\mathcal{H}_g$ by the usual formula $MZ = (AZ + B)(CZ + D)^{-1}$. The quotient $\mathcal{H}_g/\Gamma_g[q]$ is, by a well-known theorem of Baily, a quasi projective variety. We denote by $\overline{\mathcal{H}_g/\Gamma_g[q]}$ its Satake compactification. This is related to the algebra of modular forms

$$A(\Gamma_g[q]) = \bigoplus_{r=0}^{\infty} [\Gamma_g[q], r],$$

where $[\Gamma_g[q], r]$ denotes the vector space of modular forms of weight r . These are holomorphic functions $f : \mathcal{H}_g \rightarrow \mathbb{C}$ with the transformation property $f(MZ) = \det(CZ + D)^r f(Z)$ for all $M \in \Gamma_g[q]$ (and a regularity condition at the cusps in the case $g = 1$). Then one has

$$\overline{\mathcal{H}_g/\Gamma_g[q]} \cong \text{proj}(A(\Gamma_g[q])).$$

We denote by $\mathcal{J}_g \subset \mathcal{H}_g$ the set of all matrices which are period matrices of hyperelliptic (non-singular projective) curves. Then

$$\mathcal{I}_g[q] = \mathcal{J}_g/\Gamma_g[q]$$

is a quasi projective subvariety. The variety $\mathcal{I}_g[1]$ is the modular variety of hyperelliptic curves. The closure

$$\overline{\mathcal{I}_g[q]} \subset \overline{\mathcal{H}_g/\Gamma_g[q]}$$

is one of its compactifications we have to consider.

For $g \geq 3$ and $q \geq 2$, $\overline{\mathcal{I}_g[q]}$ is reducible. The number of reducible components of $\overline{\mathcal{I}_g[2]}$ is

$$2^{2g^2+g} \prod_{k=1}^g (1 - 2^{-2k}) / (2g + 2)!,$$

cf. [Ts] and it is the same for all even q . To describe the irreducible components we have to recall the thetanullwerte which for arbitrary g are defined by

$$\vartheta[m] = \sum_{n \in \mathbb{Z}^g} e^{\pi i (Z[n+m'/2] + (n+m'/2)'m'')}, \quad m = \begin{pmatrix} m' \\ m'' \end{pmatrix}.$$

Here m', m'' are columns in $\mathbb{Z}^g$. Up to the sign they depend only on $m', m'' \bmod 2$. They vanish identically if and only if ${}^t m' m''$ is odd. Hence in the case $g = 3$, there are essentially 36 such thetas and 36 irreducible components of $\overline{\mathcal{I}_3[2]}$. By classical results a matrix $Z \in \mathcal{H}_3$ is in the closure of $\mathcal{J}_3$ if and only if the product of all 36 thetas vanishes. From this one deduces that $\mathcal{I}_3[2]$ has 36 irreducible components. Each of them corresponds to a so-called “even characteristic”

$$m = \begin{pmatrix} m' \\ m'' \end{pmatrix} \in \mathbb{F}_2^6, \quad {}^t m' m'' = 0,$$

and is defined as

$$\mathcal{I}_3[2]^m := \{ Z \in \mathcal{I}_3[2]; \quad \vartheta[m](Z) = 0 \}.$$

Here we remark that $\vartheta[m]^4$ is a modular form of weight 2 on $\Gamma_2[2]$. Hence the zero locus of $\vartheta[m]$ inside $\mathcal{I}_3[2]$ is well defined. We have to consider the closure

$$\overline{\mathcal{I}_3[2]}^m \subset \overline{\mathcal{H}_3/\Gamma_g[2]}.$$

The different components are permuted under the full modular group. Hence it doesn't matter, which characteristic we use. We can take for example the zero characteristic 0. The Galois group of the covering

$$\overline{\mathcal{I}_3[2]}^m \longrightarrow \overline{\mathcal{I}_3[1]}$$

is the stabilizer of m in $\mathrm{Sp}(2, \mathbb{Z}/2\mathbb{Z})$ with respect to the usual action of this group on characteristics:

$$M\{m\} := {}^t M^{-1} m + \begin{pmatrix} (A {}^t B)_0 \\ (C {}^t D)_0 \end{pmatrix} \pmod{2}.$$

Here we use the notation A_0 for the column vector built from the diagonal of a matrix A . In case of the zero characteristic the stabilizer is the subgroup of $\mathrm{Sp}(2, \mathbb{F}_2)$, which fixes the quadratic form

$$\mathbb{F}_2^6 \longrightarrow \mathbb{F}_2, \quad m \longmapsto {}^t m' m''.$$

Hence it is our orthogonal group $\mathrm{O}(\mathbb{F}_2^6)$. This is also the image of the theta group $\Gamma_{3,\vartheta}$, where

$$\Gamma_{g,\vartheta} := \{ M \in \Gamma_g; \quad (A {}^t B)_0 \equiv (C {}^t D)_0 \equiv 0 \pmod{2} \}.$$

Hence we have

$$\mathrm{O}(\mathbb{F}_2^6) = \Gamma_{3,\vartheta} / \Gamma_3[2].$$

Next we have to make use of the theory of Thomae, who computed the theta values of hyperelliptic curves. We use (slightly modified) the beautiful approach of Mumford [Mu]. Let be

$$B = \{1, \dots, 8\}, \quad U = \{1, \dots, 4\}.$$

Recall that we consider on $\mathbb{F}_2^6$ the quadratic form

$$q : \mathbb{F}_2^6 \longrightarrow \mathbb{F}_2, \quad q(m) = {}^t m' m''$$

and the associated pairing

$$(m, n) = q(m + n) + q(m) + q(n) = {}^t m' n'' + {}^t m'' n'.$$

Their multiplicative form is

$$e(m) = (-1)^{{}^t m' m''}, \quad e(m, n) = (-1)^{{}^t m' n'' + {}^t m'' n'}.$$

Recall that a characteristic m is called even if $e(m) = 1$. Hence the even characteristics are just the isotropic vectors of the quadratic space $(\mathbb{F}_2^6, q)$. There are 36 even m . Following Mumford [Mu], we construct a map

$$T \longmapsto m(T),$$

which attaches to an subset $T \subset B$ of even order an element from $\mathbb{F}_2^6$. The basic properties of this assignment are

a) $m(T_1) = m(T_2)$ if and only if $T_1 = T_2$ or T_1 is the complement of T_2 .
Especially every element of $\mathbb{F}_2^6$ is in the image of the map m .

b) With the notation

$$T_1 \circ T_2 = T_1 \cup T_2 - T_1 \cap T_2$$

one has

$$m(T_1 \circ T_2) = m(T_1) + m(T_2) + m(\emptyset).$$

c) The characteristic $m(T)$ is even if and only if the number of elements of $T \circ U$ is divisible by 4 (hence it is 0, 4 or 8).

d) $e(m(T_1))e(m(T_2))e(m(T_1 \circ T_2)) = (-1)^{\#T_1 \cap T_2}.$

As consequence we get

$$\begin{aligned} e(m(T_1), m(T_2)) &= e(m(T_1))e(m(T_2))e(m(T_1) + (m(T_2))) = \\ &= e(m(T_1))e(m(T_2))e(m(T_1 \circ T_2) + m(\emptyset)) = \\ &= e(m(T_1))e(m(T_2))e(m(T_1 \circ T_2))e(m(\emptyset))e(m(T_1 \circ T_2), m(\emptyset)) = \\ &= (-1)^{\#T_1 \cap T_2}e(m(\emptyset))e(m(T_1 \circ T_2), m(\emptyset)) \end{aligned}$$

We observe that if we set $m(\emptyset)$ equal to the zero characteristics, then we have Mumford's procedure, but this will not be our case. To be concrete we take the following explicit realization.

$\{1, 2\}$	$\{1, 3\}$	$\{1, 4\}$	$\{1, 5\}$	$\{1, 6\}$	$\{1, 7\}$	$\{1, 8\}$	$\emptyset$
1	1	0	0	1	0	1	0
1	1	0	1	0	0	0	1
1	0	1	1	0	0	1	0
1	0	1	0	0	1	0	1
0	1	1	0	0	1	1	0
0	1	1	0	1	0	0	1

The first row contains subsets of B , the columns below are the corresponding characteristics. It is easy to verify that this assignment uniquely extends to a map with the properties a)-d). We also notice that $m(U)$ is the sum of the first three,

$${}^t m(U) = (0, 0, 0, 0, 0, 0).$$

The group $\text{Sp}(3, \mathbb{Z}/2\mathbb{Z})$ acts on the set of characteristics, we recall from [Ig1] that $e(m)$ and $e(m, n)e(m, p)e(n, p)$ are invariant. The form $e(m, n)$ results to be $\text{O}(\mathbb{F}_2^6)$ invariant.

The group S_8 acts on $B = \{1, \dots, 8\}$ and hence on the pairs $\{T, B - T\}$ of subsets of even order and on U , so we get an induced action of S_8 on $\mathbb{F}_2^6$. We

observe that $m(U) = 0$ and $\#(T \circ U)$ are invariant for the action of S_8 , in fact it generates the subgroup $O(\mathbb{F}_2^6)$ of $GL(\mathbb{F}_2^6)$. Hence we have constructed now an explicit isomorphism

$$S_8 \xrightarrow{\sim} O(\mathbb{F}_2^6).$$

This isomorphism will be used in the rest of this paper. We associate to each subset $T \subset B$ of even order a monomial $D(T)$ in the ring $\mathbb{C}[X_1, \dots, X_8]$. We have to distinguish two cases:

- 1) $T \circ U \neq 4$. In this case we attach 0.
- 2) $T \circ U = 4$. In this case we attach the following monomial of degree 12

$$e(m(T), m(\emptyset))(-1)^{\#T \cap U} \Delta \prod_{\substack{i \in T \circ U \\ j \notin T \circ U}} (X_i - X_j)^{-1}, \quad \text{where} \quad \Delta = \prod_{i < j} (X_i - X_j).$$

Because of $D(T) = D(B - T)$ the monomial

$$D(m) := D(m(T)) \quad \text{for} \quad m = m(T)$$

is well defined.

From Thomae's computation of the theta values of hyperelliptic Riemann surfaces follows (s. [Mu]):

3.1 Theorem (Thomae, 1870). *Let $\mathbb{C}[\dots \vartheta[m]^4 \dots]$ be the ring generated by the fourth powers of the 36 even thetanullwerte. Then D defines a ring homomorphism*

$$D : \mathbb{C}[\dots \vartheta[m]^4 \dots] \longrightarrow \mathbb{C}[\dots X_i - X_j \dots].$$

This homomorphism is equivariant with the standard action of $O(\mathbb{F}_2^6)$ on the thetanullwerte and the action of S_8 on the variables X_i .

The only element of the generators which lies in the kernel is $\vartheta[0]^4$. But this does of course not mean that the kernel is generated by this element.

We investigate the image. Let m be the characteristic coming from the empty set, i.e. ${}^t m = (0, 1, 0, 1, 0, 1)$, then we have

$$\begin{aligned} D(\vartheta[m]^4) &= (X_1 - X_2)(X_1 - X_3)(X_1 - X_4)(X_2 - X_3)(X_2 - X_4)(X_3 - X_4) \\ &\quad (X_5 - X_6)(X_5 - X_7)(X_5 - X_8)(X_6 - X_7)(X_6 - X_8)(X_7 - X_8). \end{aligned}$$

This is the product of the three Specht polynomials with respect to the tableaux

$$\begin{pmatrix} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 5 & 6 \\ 3 & 4 & 7 & 8 \end{pmatrix} \quad \begin{pmatrix} 1 & 2 & 5 & 6 \\ 4 & 3 & 8 & 7 \end{pmatrix}$$

This homomorphism can be given explicitly as

3.2 Proposition. *There is a homomorphism*

$$D : \mathbb{C}[\dots \vartheta[m]^4 \dots] \longrightarrow \mathbb{C}[X_1, \dots, X_8]$$

$$\searrow \qquad \qquad \qquad \uparrow$$

$$\qquad \qquad \qquad \mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}$$

Its kernel contains $\vartheta[0]^4$. An explicit description, using the notation $W_{ij} = X_i - X_j$, is given in the following table.

(0, 0, 0, 0, 0, 0)	0
(0, 0, 0, 0, 0, 1)	$- W_{13}W_{16}W_{17}W_{24}W_{25}W_{28}W_{36}W_{37}W_{45}W_{48}W_{58}W_{67}$
(0, 0, 0, 0, 1, 0)	$- W_{12}W_{13}W_{15}W_{23}W_{25}W_{35}W_{46}W_{47}W_{48}W_{67}W_{68}W_{78}$
(0, 0, 0, 0, 1, 1)	$W_{13}W_{14}W_{18}W_{25}W_{26}W_{27}W_{34}W_{38}W_{48}W_{56}W_{57}W_{67}$
(0, 0, 0, 1, 0, 0)	$- W_{14}W_{15}W_{17}W_{23}W_{26}W_{28}W_{36}W_{38}W_{45}W_{47}W_{57}W_{68}$
(0, 0, 0, 1, 0, 1)	$W_{12}W_{17}W_{18}W_{27}W_{28}W_{34}W_{35}W_{36}W_{45}W_{46}W_{56}W_{78}$
(0, 0, 0, 1, 1, 0)	$W_{15}W_{16}W_{18}W_{23}W_{24}W_{27}W_{34}W_{37}W_{47}W_{56}W_{58}W_{68}$
(0, 0, 0, 1, 1, 1)	$- W_{12}W_{14}W_{16}W_{24}W_{26}W_{35}W_{37}W_{38}W_{46}W_{57}W_{58}W_{78}$
(0, 0, 1, 0, 0, 0)	$W_{13}W_{14}W_{16}W_{25}W_{27}W_{28}W_{34}W_{36}W_{46}W_{57}W_{58}W_{78}$
(0, 0, 1, 0, 1, 0)	$- W_{13}W_{17}W_{18}W_{24}W_{25}W_{26}W_{37}W_{38}W_{45}W_{46}W_{56}W_{78}$
(0, 0, 1, 1, 0, 0)	$- W_{12}W_{14}W_{18}W_{24}W_{28}W_{35}W_{36}W_{37}W_{48}W_{56}W_{57}W_{67}$
(0, 0, 1, 1, 1, 0)	$W_{12}W_{16}W_{17}W_{26}W_{27}W_{34}W_{35}W_{38}W_{45}W_{48}W_{58}W_{67}$
(0, 1, 0, 0, 0, 0)	$W_{12}W_{15}W_{16}W_{25}W_{26}W_{34}W_{37}W_{38}W_{47}W_{48}W_{56}W_{78}$
(0, 1, 0, 0, 0, 1)	$- W_{14}W_{16}W_{18}W_{23}W_{25}W_{27}W_{35}W_{37}W_{46}W_{48}W_{57}W_{68}$
(0, 1, 0, 1, 0, 0)	$- W_{13}W_{15}W_{18}W_{24}W_{26}W_{27}W_{35}W_{38}W_{46}W_{47}W_{58}W_{67}$
(0, 1, 0, 1, 0, 1)	$W_{12}W_{13}W_{14}W_{23}W_{24}W_{34}W_{56}W_{57}W_{58}W_{67}W_{68}W_{78}$
(0, 1, 1, 0, 0, 0)	$W_{16}W_{17}W_{18}W_{23}W_{24}W_{25}W_{34}W_{35}W_{45}W_{67}W_{68}W_{78}$
(0, 1, 1, 0, 1, 1)	$W_{12}W_{15}W_{18}W_{25}W_{28}W_{34}W_{36}W_{37}W_{46}W_{47}W_{58}W_{67}$
(0, 1, 1, 1, 0, 0)	$- W_{12}W_{13}W_{17}W_{23}W_{27}W_{37}W_{45}W_{46}W_{48}W_{56}W_{58}W_{68}$
(0, 1, 1, 1, 1, 1)	$- W_{13}W_{15}W_{16}W_{24}W_{27}W_{28}W_{35}W_{36}W_{47}W_{48}W_{56}W_{78}$
(1, 0, 0, 0, 0, 0)	$W_{12}W_{14}W_{17}W_{24}W_{27}W_{35}W_{36}W_{38}W_{47}W_{56}W_{58}W_{68}$
(1, 0, 0, 0, 0, 1)	$- W_{15}W_{17}W_{18}W_{23}W_{24}W_{26}W_{34}W_{36}W_{46}W_{57}W_{58}W_{78}$
(1, 0, 0, 0, 1, 0)	$- W_{12}W_{16}W_{18}W_{26}W_{28}W_{34}W_{35}W_{37}W_{45}W_{47}W_{57}W_{68}$
(1, 0, 0, 0, 1, 1)	$W_{14}W_{15}W_{16}W_{23}W_{27}W_{28}W_{37}W_{38}W_{45}W_{46}W_{56}W_{78}$
(1, 0, 1, 0, 0, 0)	$W_{14}W_{15}W_{18}W_{23}W_{26}W_{27}W_{36}W_{37}W_{45}W_{48}W_{58}W_{67}$
(1, 0, 1, 0, 1, 0)	$- W_{15}W_{16}W_{17}W_{23}W_{24}W_{28}W_{34}W_{38}W_{48}W_{56}W_{57}W_{67}$
(1, 0, 1, 1, 0, 1)	$W_{13}W_{16}W_{18}W_{24}W_{25}W_{27}W_{36}W_{38}W_{45}W_{47}W_{57}W_{68}$
(1, 0, 1, 1, 1, 1)	$- W_{13}W_{14}W_{17}W_{25}W_{26}W_{28}W_{34}W_{37}W_{47}W_{56}W_{58}W_{68}$
(1, 1, 0, 0, 0, 0)	$W_{12}W_{13}W_{18}W_{23}W_{28}W_{38}W_{45}W_{46}W_{47}W_{56}W_{57}W_{67}$
(1, 1, 0, 0, 0, 1)	$- W_{13}W_{14}W_{15}W_{26}W_{27}W_{28}W_{34}W_{35}W_{45}W_{67}W_{68}W_{78}$
(1, 1, 0, 1, 1, 0)	$W_{14}W_{17}W_{18}W_{23}W_{25}W_{26}W_{35}W_{36}W_{47}W_{48}W_{56}W_{78}$
(1, 1, 0, 1, 1, 1)	$- W_{12}W_{15}W_{17}W_{25}W_{27}W_{34}W_{36}W_{38}W_{46}W_{48}W_{57}W_{68}$
(1, 1, 1, 0, 0, 0)	$W_{13}W_{15}W_{17}W_{24}W_{26}W_{28}W_{35}W_{37}W_{46}W_{48}W_{57}W_{68}$
(1, 1, 1, 0, 1, 1)	$W_{12}W_{13}W_{16}W_{23}W_{26}W_{36}W_{45}W_{47}W_{48}W_{57}W_{58}W_{78}$
(1, 1, 1, 1, 0, 1)	$W_{14}W_{16}W_{17}W_{23}W_{25}W_{28}W_{35}W_{38}W_{46}W_{47}W_{58}W_{67}$
(1, 1, 1, 1, 1, 0)	$W_{12}W_{14}W_{15}W_{24}W_{25}W_{36}W_{37}W_{38}W_{45}W_{67}W_{68}W_{78}$

3.3 Remark. *The variable W_{ij} appears in the monomial $D(\vartheta[m]^4)$, $m \neq 0$, if and only if m is orthogonal to the (odd) characteristic $m(T_{ij})$, where T_{ij} is defined by*

$$T_{ij} \circ U = \{i, j\}.$$

It is easy to verify this for one m by direct computation. The general case follows then because $O(\mathbb{F}_2^6)$ acts transitively on the set of non-zero even characteristics. $\square$

Finally we mention an observation of Igusa [Ig2]:

3.4 Remark. *D extends to a ring homomorphism*

$$D : A(\Gamma_3[2]) \longrightarrow \mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}.$$

An element f belongs to the kernel if and only if it vanishes on $\mathcal{I}_3[2]^0$.

Recall that on the right hand side we have the ring of Specht polynomials from section 1, which also can be identified with the ring of ball modular forms $\mathcal{R}_3/\mathcal{I}_3$ from section 2.

4. A structure theorem

Runge [Ru2] has described generators for the ring of Siegel modular forms of genus three and level two with respect to the trivial character. He gave explicitly 15 generators in weight 2 and 15 generators in weight 4. He expressed them as explicit polynomials in the 8 theta constants of second kind

$$f_a(Z) = \sum_{g \in \mathbb{Z}^3} e^{2\pi i Z[g+a/2]}, \quad a \in (\mathbb{Z}/2\mathbb{Z})^3.$$

We recall the relation to the $\vartheta[m]$:

$$\vartheta^2 \begin{bmatrix} m' \\ m'' \end{bmatrix} = \sum_{a \in \mathbb{F}_2^3} (-1)^{t_{m'm''}} f_{m'+a} f_a, \quad f_a f_b = \frac{1}{8} \sum_{c \in \mathbb{F}_2^3} (-1)^{t_{ac}} \vartheta^2 \begin{bmatrix} a+b \\ c \end{bmatrix}.$$

The classical Schottky relation R can be written as a polynomial in degree 16 of the f_a . Runge proved the basic result that this is a defining relation in the ring $\mathbb{C}[f_a]$ and moreover that this ring is normal. This implies that the subring $\mathbb{C}[f_a f_b]$ of forms of integral weight is the full ring of modular forms of integral weight and trivial character with respect to Runge's group $\Gamma_3^*[2, 4]$. This is the unique subgroup of Igusa's group $\Gamma_3[2, 4]$ which does not contain the

negative unit matrix. Recall that the Igusa group $\Gamma_g[q, 2q]$ is the subgroup of all $M \in \Gamma_g[q]$ such that $A^t B/q$ and $C^t D/q$ have even diagonal. Hence the theta group is $\Gamma_{g,\vartheta} = \Gamma_g[1, 2]$.

We recall briefly the way how Runge describes the action of the modular group on the f_a . He introduces the following group H_3 of 8×8 -matrices (indexed by the 8 elements from $\mathbb{F}_2^3$). The first generator is

$$\tilde{S} = \left(\frac{1+i}{2} \right)^3 ((-1)^{tab}).$$

To every integral diagonal 3×3 -matrices one attaches the diagonal matrix $\tilde{T}_S$ with diagonal entries $i^{S[a]}$. The group H_3 is generated by $\tilde{I}$ and by the $\tilde{T}_S$. The negative unit matrix is contained in H_3 . If we map I to $\tilde{I}$ and T_S to $\tilde{T}_S$ we obtain a homomorphism

$$\Gamma_3 \longrightarrow H_3 / \pm.$$

the kernel of this homomorphism is $\Gamma^*[2, 4]$. Moreover there is a surjective homomorphism

$$H_3 \longrightarrow \mathrm{Sp}(3, \mathbb{F}_2),$$

which sends $\tilde{I}$ and $\tilde{T}_S$ to the cosets of I and T_S . The kernel of this homomorphism is

$$N_3 := \langle iE, \tilde{T}_{2S}, \tilde{I}^{-1} \tilde{T}_{2S} I, \quad (S \text{ integral}) \rangle,$$

as correctly stated in Runge's first paper [Ru1]. In the second part of his paper [Ru2] Runge claims that N_3 agrees with

$$N'_3 := \langle \tilde{T}_{2S}, \tilde{I}^{-1} \tilde{T}_{2S} I, \quad (S \text{ integral}) \rangle.$$

But this is not the case, since $N'_3 \subset N_3$ is a subgroup of index two which doesn't contain iE . It contains only $-E$.

Now we modify Runge's results on modular forms of level two introducing the theta character v_ϑ . This is the character of the modular form $\vartheta[0]^2$, which lives on the theta group $\Gamma_{3,\vartheta} = \Gamma_3[1, 2]$. This group contains the principal congruence group of level two $\Gamma_3[2]$. We consider the ring

$$A(\Gamma_3[2]) := \bigoplus_{r \in \mathbb{Z}} [\Gamma_3[2], r, v_\vartheta^r].$$

We can get now a formal description of this ring:

4.1 Remark. *let F_a be a formal variable for $a \in \mathbb{F}_2^3$. We consider the polynomial ring $\mathbb{C}[F_a]$ and the polynomial R of degree 16 which describes the Schottky relation. Since R is invariant under H_3 the group N'_3 acts on $\mathbb{C}[F_a]/(R)$. There is a natural isomorphism*

$$(\mathbb{C}[F_a]/(R))^{N'_3} = (\mathbb{C}[F_a]^{N'_3})/(R) \xrightarrow{\sim} A(\Gamma_3[2]),$$

which replaces F_a by f_a .

From Runge's results one can derive:

4.2 Proposition. *The ring $A(\Gamma_3[2])$ is generated by*

- 1 form of weight 1 (namely $\vartheta[0]^2$), which belongs to the theta group,*
- 14 forms of weight 2,*
- 14 forms of weight 3,*
- 1 form of weight 4, which belongs to the theta group.*

The Hilbert function of the ring is $(1 - z^8)$ times

$$\frac{1 - 3z + 13z^2 - 17z^3 + 44z^4 - 17z^5 + 13z^6 - 3z^7 + z^8}{(1 - z^2)^4(1 - z)^4},$$

which is

$$1 + z + 15z^2 + 29z^3 + 135z^4 + 310z^5 + 870z^6 + 1830z^7 + 3992z^8 + 7534z^9 + 14142z^{10} + \dots$$

5. The ring of hyperelliptic modular forms of genus 3

The factor group $\Gamma_3/\Gamma_3[2]$ is isomorphic to $\mathrm{Sp}(3, \mathbb{Z}/2\mathbb{Z})$. This group contains naturally the orthogonal group $\mathrm{O}(\mathbb{F}_2^6)$ (quadratic form $m_1m_4 + m_2m_5 + m_3m_6$). Recall that the inverse image of $\mathrm{O}(\mathbb{F}_2^6)$ in the Siegel modular group is the theta group $\Gamma_{3,\vartheta}$. The group $\mathrm{O}(\mathbb{F}_2^6)$ has a simple subgroup of index two, which we denote by $\mathrm{SO}(\mathbb{F}_2^6)$. Its inverse image is a subgroup of index two $\Gamma'_{3,\vartheta} \subset \Gamma_{3,\vartheta}$. It is important to find a natural system of generators of the space $[\Gamma_3[2], 2]$. Actually we will construct 15 forms $\Theta_1, \dots, \Theta_{15}$, which generate this space and which have the property that they are permuted under the action of the group $\Gamma'_{3,\vartheta}$. For their construction we use the 36 theta constants of first kind $\vartheta[m]$.

We have thirty maximal totally isotropic spaces contained in $\mathbb{F}_2^6$. For such a subspace A we set

$$\Theta = \sum_{i=1}^7 \vartheta[m_i]^4,$$

where $m_1, \dots, m_7$ are the non-zero elements of A . The orthogonal group permutes the totally isotropic spaces, the simple group $\mathrm{SO}(\mathbb{F}_2^6)$ describes two orbits. (Two spaces are in the same orbit if and only if their intersection has even dimension.) We give the list of the subspaces in one of the two orbits. We give their entries different from 0 ordered lexicographically, i.e. m is replaced by the digit $2^5m_1 + 2^4m_2 + 2^3m_3 + 2^2m_4 + 2m_5 + m_6$. The fifteen orbits are:

$$\begin{aligned}
A_1 &= [6, 8, 14, 48, 54, 56, 62] & A_2 &= [6, 10, 12, 49, 55, 59, 61] \\
A_3 &= [7, 24, 31, 40, 47, 48, 55] & A_4 &= [7, 27, 28, 42, 45, 49, 54] \\
A_5 &= [4, 8, 12, 16, 20, 24, 28] & A_6 &= [4, 10, 14, 17, 21, 27, 31] \\
A_7 &= [5, 16, 21, 40, 45, 56, 61] & A_8 &= [5, 17, 20, 42, 47, 59, 62] \\
A_9 &= [2, 8, 10, 32, 34, 40, 42] & A_{10} &= [2, 12, 14, 33, 35, 45, 47] \\
A_{11} &= [3, 24, 27, 32, 35, 56, 59] & A_{12} &= [3, 28, 31, 33, 34, 61, 62] \\
A_{13} &= [1, 16, 17, 32, 33, 48, 49] & A_{14} &= [1, 20, 21, 34, 35, 54, 55] \\
A_{15} &= [1, 2, 3, 4, 5, 6, 7]
\end{aligned}$$

For each A_i we denote by Θ_i the corresponding sum of fourth powers of the thetanullwerte. For example with these notations we have

$$\begin{aligned}
\Theta_1 = & \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}^4 + \vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \end{bmatrix}^4 + \vartheta \begin{bmatrix} 0 & 1 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}^4 \\
& + \vartheta \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 0 \end{bmatrix}^4 + \vartheta \begin{bmatrix} 1 & 1 \\ 1 & 1 \\ 0 & 0 \end{bmatrix}^4 + \vartheta \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 0 \end{bmatrix}^4 + \vartheta \begin{bmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 0 \end{bmatrix}^4.
\end{aligned}$$

5.1 Proposition. *The forms $\Theta_1, \dots, \Theta_{15}$ give a basis of the space $[\Gamma_3[2], 2]$. As a consequence of the quartic Riemann relations the form*

$$\Theta_1 + \dots + \Theta_{15}$$

vanishes on the hyperelliptic component defined by $\vartheta[0] = 0$.

We are interested in the restriction of the ring $A(\Gamma_3[2])$ to one of the hyperelliptic components. We take the component defined by the zero locus of $\vartheta[0]$. It turns out that not only $\vartheta[0]^2$ but also the other $\Gamma_{3,\vartheta}$ invariant form in 4.2 of weight 4 vanishes on this component. We give them explicitly as polynomials in the F_a . We order them lexicographically, i.e. a is replaced by the digit $4a_1 + 2a_2 + a_3$.

$$P := F_0^2 + F_1^2 + F_2^2 + F_3^2 + F_4^2 + F_5^2 + F_6^2 + F_7^2,$$

$$\begin{aligned}
Q := & F_2^8 + 4F_4^4 F_2^2 F_3^2 + F_1^8 - 2F_7^4 F_0^4 + 4F_6^2 F_7^2 F_2^4 - 2F_7^4 F_1^4 + 24F_6^2 F_7^2 F_2^2 F_3^2 - 8F_0^2 F_2^2 F_5^2 F_6^2 - \\
& 4F_5^6 F_4^2 - 2F_3^4 F_4^4 + F_5^8 + 4F_1^4 F_2^2 F_3^2 - 2F_1^4 F_2^4 - 2F_2^4 F_0^4 - 2F_0^4 F_3^4 - 10F_1^4 F_0^4 - 4F_6^6 F_0^2 + \\
& 4F_0^4 F_3^2 F_2^2 - 2F_6^4 F_0^4 - 2F_3^4 F_1^4 + 4F_3^4 F_1^2 F_0^2 + 4F_5^4 F_2^2 F_3^2 - 2F_5^4 F_1^4 + 4F_5^4 F_0^2 F_1^2 + 4F_6^4 F_3^2 F_2^2 - \\
& 2F_7^4 F_2^4 - 2F_7^4 F_3^4 - 2F_6^4 F_1^4 - 2F_6^4 F_3^4 - 2F_6^4 F_2^4 - 2F_5^4 F_0^4 - 2F_5^4 F_2^4 - 2F_5^4 F_3^4 + 4F_7^4 F_2^2 F_3^2 + \\
& 4F_0^2 F_1^2 F_2^4 + 4F_6^4 F_1^2 F_0^2 + 4F_7^4 F_0^2 F_1^2 - 128F_1 F_0 F_3 F_2 F_5 F_4 F_7 F_6 - 10F_4^4 F_5^4 - 4F_6^6 F_7^2 - \\
& 10F_6^4 F_7^4 - 4F_6^2 F_7^6 + 4F_4^2 F_6^4 F_5^2 + 4F_4^4 F_7^2 F_6^2 + 4F_5^4 F_6^2 F_7^2 - 2F_4^4 F_6^4 - 2F_4^4 F_7^4 - 2F_5^4 F_6^4 + \\
& F_6^8 + F_4^8 + F_7^8 - 2F_5^4 F_7^4 - 4F_4^4 F_5^2 + F_0^8 + F_3^8 + 4F_0^2 F_1^2 F_4^4 + 4F_4^4 F_5^2 F_7^4 - 2F_2^4 F_4^4 - 4F_0^6 F_1^2 - \\
& 4F_2^6 F_3^2 - 10F_2^4 F_3^4 - 4F_2^2 F_3^6 + 4F_2^4 F_4^2 F_5^2 + 4F_1^4 F_6^2 F_7^2 + 4F_1^4 F_4^2 F_5^2 + 4F_0^4 F_6^2 F_7^2 + 4F_3^4 F_6^2 F_7^2 + \\
& 4F_0^4 F_4^2 F_5^2 - 2F_0^4 F_4^4 - 2F_1^4 F_4^4 + 4F_3^4 F_4^2 F_5^2 + 24F_0^2 F_1^2 F_2^2 F_3^2 + 8F_0^2 F_2^2 F_7^2 F_5^2 - 8F_0^2 F_2^2 F_7^2 F_4^2 + \\
& 8F_1^2 F_3^2 F_7^2 F_5^2 - 8F_1^2 F_3^2 F_7^2 F_4^2 - 8F_1^2 F_3^2 F_6^2 F_5^2 + 8F_4^2 F_6^2 F_2^2 F_0^2 + 8F_4^2 F_6^2 F_1^2 F_3^2 - \\
& 8F_4^2 F_6^2 F_1^2 F_2^2 + 8F_1^2 F_2^2 F_4^2 F_7^2 - 8F_0^2 F_3^2 F_5^2 F_7^2 + 8F_0^2 F_3^2 F_5^2 F_6^2 + 8F_0^2 F_3^2 F_4^2 F_7^2 - \\
& 8F_0^2 F_3^2 F_4^2 F_6^2 + 8F_1^2 F_2^2 F_5^2 F_6^2 - 8F_1^2 F_2^2 F_5^2 F_7^2 + 24F_0^2 F_1^2 F_4^2 F_5^2 + 24F_4^2 F_5^2 F_7^2 F_6^2 + \\
& 24F_6^2 F_7^2 F_1^2 F_0^2 + 24F_4^2 F_5^2 F_2^2 F_3^2
\end{aligned}$$

5.2 Proposition. *The ring*

$$\mathbb{C}[F_a]/(P, Q)$$

is normal of Krull dimension 6.

Proof. Since it is an complete intersection, one has only to show the the codimension of the singular locus is at least two. This can be checked with a computer. $\square$

We need the invariant ring

$$B(\Gamma_3[2]) := (\mathbb{C}[F_a]/(P, Q))^{N'_3} = \mathbb{C}[F_a]^{N'_3}/(P, Q).$$

The zero locus of $\vartheta[0]$ is an irreducible subset of $\mathcal{H}_3/\Gamma_3[2]$. The restriction of the ring of modular forms $A(\Gamma_3[2])$ to this hyperelliptic component hence is

$$A(\Gamma_3[2])/\text{rad}((\vartheta[0]^2))$$

5.3 Theorem. *The restriction of the ring of modular forms $A(\Gamma_3[2])$ to the hyperelliptic component, set theoretically defined by $\vartheta[0] = 0$, is*

$$A(\Gamma_3[2])/\text{rad}((\vartheta[0]^2)) \cong B(\Gamma_3[2]) = \mathbb{C}[F_a]^{N'_3}/(P, Q).$$

It is generated by 14 forms of weight 2 and 14 forms of weight 3. The Hilbert function of this ring is the product of the Hilbert function of $A(\Gamma_3[2])$ (see 4.2) with $(1 - z)(1 - z^4)/(1 - z^8)$,

$$1 + 14z^2 + 14z^3 + 105z^4 + 175z^5 + 546z^6 + 946z^7 + 2057z^8 \dots$$

For the proof we have to describe the relations between the generators. For this we need a method which allows to decide whether a system $f_1, \dots, f_m$ of homogenous polynomials in the variables F_a is linear independent in the quotient $\mathbb{C}[F_a]/(P, Q)$. Of course we assume that all f_i have the same degree. We used the following method: First of all, we computed a Groebner basis of the ideal (P, Q) , using the computer algebra SINGULAR. Then we computed the *normal forms* $g_1, \dots, g_m$ of the f_i with respect to this basis. The point now is that the $f_1, \dots, f_m$ are linearly independent in the factor ring $\mathbb{C}[F_a]/(P, Q)$ if and only if the $g_1, \dots, g_m$ are linearly independent in the polynomial ring $\mathbb{C}[F_a]$. This is of course a problem in linear algebra. Using this method we proved:

5.4 Proposition. *The space of forms in $B(\Gamma_3[2])$ of weight four, which is generated by products of two forms of weight two has dimension 105. The space of forms of weight six, which is generated by products of three forms of weight two has dimension 546.*

Corollary 1. *The 14 forms of weight two in generate in $B(\Gamma_3[2])$ all forms of weight 4 and 6.*

Corollary 2. *There are 560 monomials of degree 4 in the 14 generators of weight two. Hence there is a 14-dimensional space of cubic relations.*

It seems to be natural now to consider the subring

$$B'(\Gamma_3[2]) \subset B(\Gamma_3[2])$$

generated by the 14 elements of weight 2. Because of 5.4 one can conjecture that both rings agree in even weights. We will see that this is true (5.8).

Several times a 14-dimensional space occurred. This depends on the fact that the group $O(6, \mathbb{F}_2) \cong S_8$ admits an irreducible irreducible 14-dimensional representation. There are actually to isomorphism classes which are intertwined by the signum character. All 14-dimensional spaces, which occurred so far, are irreducible. Next we describe the 14 cubic relations:

5.5 Lemma. *Restricting to the hyperelliptic component, the following ternary polynomial is contained in the ideal generated by the quartic Riemann relations:*

$$\begin{aligned} & 18 \cdot \Theta_6 \cdot \Theta_{11} \cdot \Theta_{14} + 18 \cdot \Theta_8 \cdot \Theta_9 \cdot \Theta_{14} + 18 \cdot \Theta_2 \cdot \Theta_{13} \cdot \Theta_{14} + 9 \cdot \Theta_1 \cdot \Theta_{12}^2 + 9 \cdot \Theta_1 \cdot \Theta_2^2 + 18 \cdot \Theta_3 \cdot \\ & \Theta_{14} \cdot \Theta_{15} - 43 \cdot \Theta_1^3 + 9 \cdot \Theta_1 \cdot \Theta_9^2 + 2 \cdot \Theta_4^3 + 2 \cdot \Theta_8^3 + 2 \cdot \Theta_{11}^3 + 2 \cdot \Theta_{13}^3 + 18 \cdot \Theta_2 \cdot \Theta_5 \cdot \Theta_6 - 18 \cdot \Theta_5^2 \cdot \\ & \Theta_9 + 2 \cdot \Theta_3^3 + 9 \cdot \Theta_1 \cdot \Theta_5^2 - 18 \cdot \Theta_7 \cdot \Theta_{11}^2 - 18 \cdot \Theta_4 \cdot \Theta_{14}^2 + 2 \cdot \Theta_{12}^3 + 18 \cdot \Theta_4 \cdot \Theta_5 \cdot \Theta_8 - 18 \cdot \Theta_2 \cdot \\ & \Theta_{15}^2 + 9 \cdot \Theta_1 \cdot \Theta_3^2 + 18 \cdot \Theta_6 \cdot \Theta_{12} \cdot \Theta_{13} + 18 \cdot \Theta_8 \cdot \Theta_{11} \cdot \Theta_{15} + 9 \cdot \Theta_1 \cdot \Theta_{15}^2 + 18 \cdot \Theta_7 \cdot \Theta_{10} \cdot \Theta_{14} + \\ & 9 \cdot \Theta_1 \cdot \Theta_8^2 + 2 \cdot \Theta_{14}^3 + 18 \cdot \Theta_2 \cdot \Theta_7 \cdot \Theta_8 + 2 \cdot \Theta_9^3 + 9 \cdot \Theta_1 \cdot \Theta_{10}^2 + 18 \cdot \Theta_3 \cdot \Theta_{10} \cdot \Theta_{12} + 9 \cdot \Theta_1 \cdot \Theta_4^2 + 2 \cdot \\ & \Theta_5^3 + 18 \cdot \Theta_3 \cdot \Theta_6 \cdot \Theta_8 - 18 \cdot \Theta_5 \cdot \Theta_9^2 + 2 \cdot \Theta_{15}^3 + 2 \cdot \Theta_6^3 + 18 \cdot \Theta_7 \cdot \Theta_9 \cdot \Theta_{13} + 18 \cdot \Theta_7 \cdot \Theta_{12} \cdot \Theta_{15} + \\ & 30 \cdot \Theta_1 \cdot \Theta_7 \cdot \Theta_{11} + 30 \cdot \Theta_1 \cdot \Theta_3 \cdot \Theta_{13} + 18 \cdot \Theta_2 \cdot \Theta_{11} \cdot \Theta_{12} + 9 \cdot \Theta_1 \cdot \Theta_{13}^2 + 30 \cdot \Theta_1 \cdot \Theta_4 \cdot \Theta_{14} + 30 \cdot \\ & \Theta_1 \cdot \Theta_2 \cdot \Theta_{15} + 18 \cdot \Theta_5 \cdot \Theta_{10} \cdot \Theta_{15} + 9 \cdot \Theta_1 \cdot \Theta_{14}^2 + 18 \cdot \Theta_4 \cdot \Theta_9 \cdot \Theta_{12} - 6 \cdot \Theta_{13} \cdot \Theta_{14} \cdot \Theta_{15} - 6 \cdot \Theta_5 \cdot \\ & \Theta_8 \cdot \Theta_{14} - 6 \cdot \Theta_3 \cdot \Theta_4 \cdot \Theta_{15} - 6 \cdot \Theta_9 \cdot \Theta_{10} \cdot \Theta_{15} - 6 \cdot \Theta_6 \cdot \Theta_7 \cdot \Theta_{14} - 6 \cdot \Theta_4 \cdot \Theta_5 \cdot \Theta_{12} - 6 \cdot \Theta_7 \cdot \Theta_8 \cdot \\ & \Theta_{15} - 6 \cdot \Theta_3 \cdot \Theta_7 \cdot \Theta_9 - 6 \cdot \Theta_9 \cdot \Theta_{12} \cdot \Theta_{14} - 6 \cdot \Theta_4 \cdot \Theta_8 \cdot \Theta_9 - 6 \cdot \Theta_2 \cdot \Theta_7 \cdot \Theta_{12} - 6 \cdot \Theta_3 \cdot \Theta_6 \cdot \Theta_{12} - \\ & 6 \cdot \Theta_4 \cdot \Theta_7 \cdot \Theta_{10} - 6 \cdot \Theta_{11} \cdot \Theta_{12} \cdot \Theta_{15} - 6 \cdot \Theta_6 \cdot \Theta_8 \cdot \Theta_{13} - 6 \cdot \Theta_5 \cdot \Theta_7 \cdot \Theta_{13} - 6 \cdot \Theta_{10} \cdot \Theta_{12} \cdot \Theta_{13} + \\ & 9 \cdot \Theta_1 \cdot \Theta_6^2 - 6 \cdot \Theta_3 \cdot \Theta_8 \cdot \Theta_{10} - 6 \cdot \Theta_9 \cdot \Theta_{11} \cdot \Theta_{13} - 6 \cdot \Theta_2 \cdot \Theta_4 \cdot \Theta_{13} - 6 \cdot \Theta_5 \cdot \Theta_6 \cdot \Theta_{15} - 6 \cdot \Theta_2 \cdot \Theta_6 \cdot \\ & \Theta_9 - 6 \cdot \Theta_2 \cdot \Theta_5 \cdot \Theta_{10} - 6 \cdot \Theta_4 \cdot \Theta_6 \cdot \Theta_{11} - 6 \cdot \Theta_{10} \cdot \Theta_{11} \cdot \Theta_{14} - 6 \cdot \Theta_2 \cdot \Theta_8 \cdot \Theta_{11} - 6 \cdot \Theta_2 \cdot \Theta_3 \cdot \Theta_{14} + \\ & 18 \cdot \Theta_5 \cdot \Theta_{11} \cdot \Theta_{13} + 18 \cdot \Theta_2 \cdot \Theta_9 \cdot \Theta_{10} + 18 \cdot \Theta_4 \cdot \Theta_6 \cdot \Theta_7 + 18 \cdot \Theta_6 \cdot \Theta_9 \cdot \Theta_{15} + 18 \cdot \Theta_4 \cdot \Theta_{13} \cdot \Theta_{15} + \\ & 18 \cdot \Theta_8 \cdot \Theta_{10} \cdot \Theta_{13} + 18 \cdot \Theta_2 \cdot \Theta_3 \cdot \Theta_4 + 18 \cdot \Theta_4 \cdot \Theta_{10} \cdot \Theta_{11} + 9 \cdot \Theta_1 \cdot \Theta_{11}^2 + 18 \cdot \Theta_3 \cdot \Theta_9 \cdot \Theta_{11} - 18 \cdot \\ & \Theta_2^2 \cdot \Theta_{15} - 18 \cdot \Theta_6 \cdot \Theta_{10}^2 - 18 \cdot \Theta_6^2 \cdot \Theta_{10} - 18 \cdot \Theta_3^2 \cdot \Theta_{13} - 18 \cdot \Theta_3 \cdot \Theta_{13}^2 - 18 \cdot \Theta_8 \cdot \Theta_{12}^2 - 18 \cdot \Theta_8^2 \cdot \\ & \Theta_{12} - 18 \cdot \Theta_4^2 \cdot \Theta_{14} + 2 \cdot \Theta_{10}^3 + 2 \cdot \Theta_7^3 + 30 \cdot \Theta_1 \cdot \Theta_8 \cdot \Theta_{12} + 30 \cdot \Theta_1 \cdot \Theta_5 \cdot \Theta_9 + 2 \cdot \Theta_2^3 + 30 \cdot \Theta_1 \cdot \\ & \Theta_6 \cdot \Theta_{10} - 6 \cdot \Theta_3 \cdot \Theta_5 \cdot \Theta_{11} + 18 \cdot \Theta_3 \cdot \Theta_5 \cdot \Theta_7 - 18 \cdot \Theta_7^2 \cdot \Theta_{11} + 9 \cdot \Theta_1 \cdot \Theta_7^2 + 18 \cdot \Theta_5 \cdot \Theta_{12} \cdot \Theta_{14} \end{aligned}$$

Its orbit under the group $\Gamma_{3,\theta'}$ gives a system of 14 relations.

The relations of degree ≤ 3 above do not generate the ideal of all relations. We also need some quartic relations. To compute them one needs the expressions of the $\vartheta[m]^4$ as linear combinations of the $\Theta_1, \dots, \Theta_{15}$. Here is one:

$$6\vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}^4 = \Theta_{13} + \Theta_{14} + \Theta_{15}.$$

(We use the notation $\vartheta[m] = \vartheta[m', m'']$.) The others are obtained by applying $O(\mathbb{F}_2^6)$. Actually there will be two types of quartic relations between the $\vartheta[m]^4$. Let M be a set of characteristics. We denote by $\vartheta[M]$ the product of all $\vartheta[m]$ with $m \in M$. Let M now be a two dimensional $\mathbb{F}_2$ -vector space of characteristics. There are three cosets $M, a+M, b+M$ containing only even characteristics. We have the Riemann relation

$$\vartheta[M] = \vartheta[M+a] \pm \vartheta[M+b].$$

If we set $\vartheta[0]$ to zero we obtain the relation $\vartheta[a+M]^4 = \vartheta[b+M]^4$.

5.6 Proposition. *There are 210 relations between the Θ_i (considered on the hyperelliptic component $\vartheta[0](\tau) = 0$) which come from the relations*

$$\vartheta[a+M]^4 = \vartheta[b+M]^4.$$

Here M is a two-dimensional $\mathbb{F}_2$ -vector space of characteristics and $a+M$ and $b+M$ are the two orbits consisting only of even characteristics.

One can expand these relations as quartic polynomials in the Θ_i . We don't print them here.

We also need quartic relations which contain only squares of thetas : They are of the form

$$\vartheta[m_1]^2 \vartheta[m_2]^2 \pm \vartheta[m_3]^2 \vartheta[m_4]^2 = \vartheta[m_5]^2 \vartheta[m_6]^2 \pm \vartheta[m_7]^2 \vartheta[m_8]^2.$$

Now assuming $\vartheta[m_1] = \vartheta[0] = 0$ we have

$$\pm \vartheta[m_3]^2 \vartheta[m_4]^2 = \vartheta[m_5]^2 \vartheta[m_6]^2 \pm \vartheta[m_7]^2 \vartheta[m_8]^2.$$

Squaring we get

$$\vartheta[m_3]^4 \vartheta[m_4]^4 - \vartheta[m_5]^4 \vartheta[m_6]^4 - \vartheta[m_7]^4 \vartheta[m_8]^4 = \pm 2\vartheta[m_5]^2 \vartheta[m_6]^2 \vartheta[m_7]^2 \vartheta[m_8]^2 :$$

Squaring again we get a quartic relation among $\vartheta[m]^4$. This can be expressed in the Θ_i . The total number of these relations between the Θ_i is 105.

5.7 Proposition. *The relations of the type (considered in $B(\Gamma_3[2])$)*

$$(\vartheta[m_3]^4\vartheta[m_4]^4 - \vartheta[m_5]^4\vartheta[m_6]^4 - \vartheta[m_7]^4\vartheta[m_8]^4)^2 = 4\vartheta[m_5]^4\vartheta[m_6]^4\vartheta[m_7]^4\vartheta[m_8]^4$$

give 105 relations between the Θ_i .

We will not give the explicit polynomials in the Θ_i . The relations described so far are still not all. The Schottky relation

$$\left(\sum_m \vartheta[m]^8\right)^2 - 8 \sum_m \vartheta[m]^{16}$$

is an extra relation. The expansion of this relation as polynomial in the Θ_i is very big. We don't print it.

Remember that we now have the following system of relations between the 15 functions Θ_i considered in $B(\Gamma_3[2])$.

- 1) One linear relation,
- 2) 14 cubic relations,
- 3) two systems of quartic relations one consisting of 105 the other of 210 relations,
- 4) an extra quartic relation.

This set of relations is permuted under the action of $\Gamma'_{3,\vartheta}$. We consider the ideal $\mathcal{R}$ generated by all these relations.

Using the computer algebra system **SINGULAR** one can get a Gröbner basis of this ideal, which allows to do several computations, for example it is possible to get the Hilbert function of the algebra

$$\mathbb{C}[T_1, \dots, T_{15}]/\mathcal{R}.$$

Here the T_i are formal variables which stand for the Θ_i .

SINGULAR gives the following Hilbert function:

$$(1 + 8z^2 + 36z^4 + 106z^6 + 91z^8 + 14z^{10})/(1 - z^2)^6.$$

(The weight of Θ_i is two.) One checks immediately that this series agrees with the even part of the Hilbert series of the ring $B(\Gamma_3[2])$ as has been described in 6.2. Hence we obtain that our ideal describes all relations.

We mention that the our system of relations is not minimal. The system of relations has the advantage to be invariant under the group $\Gamma'_{3,\vartheta}$.

The final result is:

5.8 Theorem. *The subring $B^{(2)}(\Gamma_3[2])$ of forms of even weight of the ring*

$$B(\Gamma_3[2]) = A(\Gamma_3[2]) / (\text{rad}((\vartheta[0]^2)))$$

is generated by 15 forms $\Theta_1, \dots, \Theta_{15}$. The ideal of relations is generated by one linear relation, 14 ternary relations and 127 quartic relations: The Hilbert function is given by the formula

$$\frac{1 + 8z^2 + 36z^4 + 106z^6 + 91z^8 + 14z^{10}}{(1 - z^2)^6} = \\ 1 + 14z^2 + 105z^4 + 546z^6 + 2057z^8 + 6062z^{10} + 14945z^{12} + \\ 32306z^{14} + 63217z^{16} + 114478z^{18} + \dots$$

The projective variety of this ring is the hyperelliptic component in the Satake compactified Siegel modular variety $\overline{\mathcal{H}_3/\Gamma_3[2]}$. Hence this closure is a normal variety.

6. Blowing up

We want to investigate the homomorphism

$$B^{(2)}(\Gamma_3[2]) = \mathbb{C}[\dots \vartheta[m]^4 \dots] \longrightarrow \mathbb{C}[Y_1, \dots, Y_{14}] / \mathcal{J}.$$

Here the $\vartheta^4[m]$ are understood as forms on the hyperelliptic component. Recall that the $\vartheta^4[m]$ can be expressed by means of the Θ_i . Hence the ideal of relations between the $\vartheta^4[m]$ (restricted to the hyperelliptic component) is known from 5.8. The right hand side is the graded algebra of the ball quotient. The ideal $\mathcal{J}$ of relations has also been described explicitly in 1.1. Recall that we have an explicit isomorphism $O(\mathbb{F}_2^6) \cong S_8$. The group $O(\mathbb{F}_2^6)$ acts transitively on the even characteristics $m \neq 0$. The group S_8 acts linearly on the Y_i . (They are polynomials in 8 variables $X_1, \dots, X_8$ on which the group S_8 acts by permutation.)

Using Thomae's theorem 3.1 and the expressions of the Y_i as polynomials in the X_i (section 1) one can compute explicit expressions of the images of the $\vartheta[m]^4$. The base locus ideal is the ideal generated by the images. Since the above homomorphism is $O(\mathbb{F}_2^6) = S_8$ -equivariant, it is sufficient to give the image of one $\vartheta[m]^4$.

6.1 Lemma. *The homomorphism*

$$\mathbb{C}[\dots \vartheta[m]^4 \dots] \longrightarrow \mathbb{C}[Y_1, \dots, Y_{14}] / \mathcal{J}$$

is defined by

$$\vartheta \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}^4 \longmapsto (Y_1 - Y_{10} + Y_{11} - Y_{14})(Y_1 - Y_2 - Y_6 + Y_7 + Y_8 - Y_9 + Y_{11} - Y_{13})(Y_8 - Y_9).$$

The others are obtained applying S_8 .

The base locus is defined by the ideal which is generated by the images of the $\vartheta[m]^4$.

6.2 Proposition. *The ideal, which is generated by the images of the $\vartheta[m]^4$ in the ring $\mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J}$ is the intersection of 56 ideals which are generated by linear forms. One of them is*

$$(Y_2, Y_3, Y_5, Y_6, Y_8, Y_9, Y_{10}, Y_{11}, Y_{13}, Y_{14})$$

The others are obtained applying S_8 .

This can be verified by a SINGULAR calculation. However: in the last section we shall give a more geometric and intrinsic description of the components of zero set of the ideal generated by the images of the $\vartheta[m]^4$.

Let $\mathfrak{A}$ be an ideal in a (commutative and with unity) noetherian ring A . The blow up of A along $\mathfrak{A}$ is the graded A -algebra

$$\mathrm{Bl}(A, \mathfrak{A}) := \bigoplus_{n=0}^{\infty} \mathfrak{A}^n.$$

We can consider $\mathrm{Bl}(A, \mathfrak{A})$ as a subring of $A[T]$, using the embedding

$$\mathrm{Bl}(A, \mathfrak{A}) \longrightarrow A[T], \quad a \in \mathfrak{A}^n \longmapsto aT^n.$$

Now we assume

$$A = \mathbb{C}[X_1, \dots, X_n]/\mathfrak{a}.$$

We choose polynomials $Q_1, \dots, Q_m$ in $\mathbb{C}[X_1, \dots, X_n]$, whose images in A generate $\mathfrak{A}$. We consider the homomorphism of polynomial rings

$$\mathbb{C}[X_1, \dots, X_n, Y_1, \dots, Y_m] \longrightarrow \mathbb{C}[X_1, \dots, X_n, T], \quad Y_i \longmapsto Q_i T.$$

We denote by $\tilde{\mathfrak{A}}$ the the inverse image of the ideal generated by $\mathfrak{a}$. Then we have

$$\mathrm{Bl}(A, \mathfrak{A}) = \mathbb{C}[X_1, \dots, X_n, Y_1, \dots, Y_m]/\tilde{\mathfrak{A}}.$$

We need a modification of this construction. Assume that A already is a graded algebra and $\mathfrak{A}$ a graded ideal. Then $\mathrm{Bl}(A, \mathfrak{A})$ is a bigraded algebra and one can define in an obvious way the projective variety $\mathrm{biproj}(\mathrm{Bl}(A, \mathfrak{A}))$ together with a morphism

$$\mathrm{biproj}(\mathrm{Bl}(A, \mathfrak{A})) \longrightarrow \mathrm{proj} A.$$

We describe this map in terms of coordinates: For simplicity we assume that the degree of the X_i are one and that the Q_i all are of the same degree. The variety $\mathrm{biproj}(\mathrm{Bl}(A, \mathfrak{A}))$ then consists of all pairs $([x], [y])$, $[x] \in P^{n-1}(\mathbb{C})$,

$[y] \in P^{m-1}(\mathbb{C})$ such that $P(x, y) = 0$ for all polynomials $P(X, Y) \in \tilde{\mathfrak{A}}$ which are homogenous in $X = (X_1, \dots, X_n)$ and $Y = (Y_1, \dots, Y_m)$.

We want to apply this in the following situation: We consider the rings

$$B = \mathbb{C}[Y_1, \dots, Y_{14}]/\mathcal{J} \quad \text{and} \quad A = \mathbb{C}[\dots \vartheta[m]^4 \dots].$$

Here the $\vartheta[m]^4$ are understood to be restricted to the hyperelliptic component, hence $\vartheta[0]^4 = 0$. We recall that there is a homomorphism

$$A = \mathbb{C}[\dots \vartheta[m]^4 \dots] \longrightarrow B.$$

The ideal which we want to blow up is generated by the images of the $\vartheta[m]^4$ in B . We have shown that this ideal is generated by 56 linear forms. We denote the blow up of this ideal by B^* and the associated variety by Y^* . This is a model which lies over the Siegel model $X = \text{proj}(A)$ and over the ball-model $Y = \text{proj}(B)$.

$$\begin{array}{ccc} & Y^* & \\ \swarrow & & \searrow \\ Y & \text{-----} & X \end{array}$$

We want to get explicit information about the model Y^* . Actually it is too difficult to blow up the 56 components of the base locus in one step. But this is not necessary. To get information about the blow up of a small neighborhood of a given point in Y , it is sufficient to blow up those of the linear components which contain this point. To make use of this we need the intersection behavior of the 56. A direct computation shows:

6.3 Lemma. *When a subset of the 56 linear components has a common intersection, it intersects also in a cusp. In each cusp meet 8 of the linear components.*

For example the 8 components defined by the following ideals (each generated

by 10 linear forms)

$$\begin{aligned}
&(Y_2, Y_3, Y_5, Y_6, Y_8, Y_9, Y_{10}, Y_{11}, Y_{13}, Y_{14}), \\
&(Y_1, Y_2, Y_3 - Y_{11} + Y_{14}, Y_4 - Y_{12} + Y_{14}, Y_5, Y_6 - Y_{11}, Y_7 - Y_{12}, \\
&\quad Y_8 - Y_9, Y_{10}, -Y_{14} + Y_{13}), \\
&(Y_1, Y_2, Y_3, Y_5 - Y_{13} + Y_{14}, Y_6 - Y_{13}, Y_7 - Y_{12}, Y_8 - Y_{13}, \\
&\quad Y_9 - Y_{14}, Y_{10} - Y_{13} + Y_{14}, Y_{11} - Y_{13}), \\
&(Y_1, Y_2 - Y_{11} + Y_{13}, Y_3 - Y_{11} + Y_{13}, Y_4 - Y_{12}, Y_5 - Y_{11}, Y_6 - Y_{11}, \\
&\quad Y_7 - Y_{12}, Y_9, Y_{10} - Y_{11}, Y_{14}), \\
&(Y_1, Y_2, Y_3, Y_4 - Y_7, Y_5, Y_6, Y_{10}, Y_{11}, Y_{13}, Y_{14}), \\
&(Y_1 - Y_{10} - Y_{14}, Y_2 - Y_{10} - Y_{14}, Y_3, Y_4 - Y_9 - Y_{12} + Y_{14}, Y_5 - Y_{10}, \\
&\quad Y_6, Y_7 - Y_9 - Y_{12} + Y_{14}, Y_8, Y_{11}, Y_{13}), \\
&(Y_2 - Y_{10}, Y_3 - Y_{11}, Y_4 - Y_{12}, Y_5 - Y_{10}, Y_6 - Y_{11}, Y_7 - Y_{12}, \\
&\quad Y_8, Y_9, Y_{13}, Y_{14}), \\
&(Y_1 + Y_7 - Y_{12}, Y_2 + Y_{13} - Y_{14}, Y_3 + Y_{13}, Y_4 - Y_{12} + Y_{14}, \\
&\quad Y_5, Y_6, Y_8, Y_9, Y_{10}, Y_{11}),
\end{aligned}$$

meet in a cusp with the coordinates $[0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 0, 1, 0, 0]$.

The intersection of the 8 ideals, which are generated by these linear forms is generated by 17 elements of degree ≤ 2 . We will not print them and mention just that this ideal is simple enough to be blown up by means of SINGULAR. In this way we obtain for each cusp s a partial blow up Y_s^* . They exhaust Y^* in the following sense. Consider for each cusp the complement of the remaining $48 = 56 - 8$ components in Y_s^* . These are quasi projective varieties which give an open covering of Y^* . This means that we have a description of Y^* by explicit equations. We will not print them here but mention just some consequences which can be taken from them.

6.4 Proposition. *The model Y^* is smooth outside the inverse images of the cusps. It is not smooth everywhere. The inverse image of a cusp is a two dimensional irreducible variety.*

To desingularize one needs one further blow up:

6.5 Proposition. *The blow up of Y^* along the inverse images of the cusps is a non-singular model $\tilde{Y}$.*

For the proof it is convenient to proceed slightly different. We first blow up the cusps and then the inverse images of the 56 linear spaces. The result is the same. Direct computations show:

6.6 Proposition. *The blow up of the ball model Y along the cusps is a smooth model Y' . The inverse images of the cusps are irreducible. The intersection of three different of the strict transforms of the 56 linear subspaces of Y is empty.*

Since Y' is smooth, the inverse image of the union of the 56 linear spaces and the union of the strict transformed only differ by an invertible ideal. Since the blow-up of two ideals which differ only by an invertible ideal is the same, we now only have to blow-up the union of the 56 strict transforms. For the proof of 6.5 it now is sufficient to show that the blow up of Y' along the union of two of the 56 strict transformed is smooth. Actually the strict transformed are smooth and the local analytical behavior in an intersection point is the same as $z_1 = z_2 =$ and $z_3 = z_4 =$ in a $\mathbb{C}^5$. This shows that the blow-up remains smooth and proves 6.5.

Finally we compare the model $\tilde{Y}$ with $\bar{M}_{0,8}$. Recall that $M_{0,8}$ is the moduli space of 8 ordered points on a projective line and $\bar{M}_{0,8}$ the Mumford-compactification by marked stable curves. It is known [Ka] (s. also [AL]) that this is a projective smooth variety and that there exist regular contraction maps ([Ka], s. also [AL])

$$\bar{M}_{0,8} \longrightarrow Y.$$

From the universal property of blowing up one obtains a regular map

$$\bar{M}_{0,8} \longrightarrow \tilde{Y}.$$

We claim that this is biholomorphic.

6.7 Theorem. *The two diagrams*

$$\begin{array}{ccc} & \tilde{Y} & \\ \swarrow & & \searrow \\ Y & \text{-----} & X \end{array} \quad \begin{array}{ccc} & \bar{M}_{0,8} & \\ \swarrow & & \searrow \\ Y & \text{-----} & X \end{array}$$

are isomorphic.

The proof rests on the following simple

6.8 Lemma. *Let V, W be smooth complete varieties and $f : V \rightarrow W$ a regular map which induces a biholomorphic map $V_0 \rightarrow W_0$ of certain Zariski open subsets $V_0 \subset V$, $W_0 \subset W$. Assume that $V - V_0$ and $W - W_0$ are of pure codimension one and that the number of irreducible components in both cases is the same. Then f is biholomorphic.*

We omit the simple proof. We have to apply it to $V = \bar{M}_{0,8}$, $V_0 = M_{0,8}$, $W = \tilde{Y}$ and W_0 the complement of the inverse images of the Heegner divisors in Y .

Proof of 6.7. Firstly we recall shortly the irreducible components of $\bar{M}_{0,8} - M_{0,8}$. Their generic points are marked curves $(C, x_1, \dots, x_8)$, where C is the union of two P^1 with one intersection point (normal crossing). We have to choose a point on the first P^1 and b on the second one, $a + b = 8$. The possibilities to get a stable marked curve are $(a, b) = (6, 2), (5, 3), (4, 4)$. In the case $(a, b) = (6, 2)$ we have $28 = \binom{8}{2}$ possibilities, in the case $(a, b) = (5, 3)$ there are $56 = \binom{8}{3}$ and in the case $(a, b) = (4, 4)$ there are $35 = \binom{8}{4}/2$ possibilities. The denominator 2 comes from the fact that the role of the two P^1 can be interchanged. Hence we have 28, 56 and 35 irreducible components. The group S_8 acts on each of the three systems transitively.

This picture can be recovered in $\tilde{Y}$. We have 28 irreducible subvarieties of codimension 1 which map to the 28 Heegner divisors in Y . Hence these 28 are visible already in Y . Then we have the 56 irreducible subvarieties of codimension 1 which come from blowing up the 56 components of the base locus. These 56 are visible in Y^* . Finally we have the 35 irreducible subvarieties of codimension 1 which come from blowing up the cusps.

Since $O(\mathbb{F}_2^6)$ permutes the three systems, this counting also shows:

6.9 Proposition. *The boundary $\bar{M}_{0,8} - M_{0,8}$ is the union of $28 + 56 + 35$ irreducible subvarieties of codimension one which correspond to distributions $(6, 2), (5, 3), (4, 4)$ of eight points on two crossing P^1 . Under the map $\bar{M}_{0,8} \rightarrow Y$ type $(6, 2)$ -components map to the 28 Heegner divisors, type $(5, 3)$ -components to the 56 components of the base locus and type $(4, 4)$ -components are contracted to the cusps.*

In the next section we shall consider also the image of the boundary $\bar{M}_{0,8} - M_{0,8}$ under the map $\bar{M}_{0,8} \rightarrow X$

7. A combinatorial approach

In this section we want to give a combinatorial approach to the two different compactifications X and Y of the hyperelliptic component. We will restrict our attention to those points in X and Y that do not correspond to non singular complete hyperelliptic curves. We shall denote by X_0 and Y_0 the open set in X and Y corresponding to hyperelliptic curves. Thus we need at least a set theoretical description of the complementary loci.

Firstly we recall shortly the structure of $X - X_0$. We use the embedding of the Siegel half plane $\mathcal{H}_g$ into a Grassmanian and we denote by $\mathcal{H}_g^*$ the union of $\mathcal{H}_g$ with the rational boundary components. The Satake compactification then is $\mathcal{H}_g^*/\Gamma_g[q]$. Details can be found for example in [Fr3].

We restrict now to $g = 3$. Recall that X_0 is the subset of all points $\mathcal{H}_3/\Gamma_3[2]$, which are represented by a (smooth) hyperelliptic Riemann surface for which

$\vartheta[0]$ vanishes and that X is the closure of X_0 in the Satake compactification. We recall that a point $Z \in \mathcal{H}_3$ is reducible if it is conjugate with respect to the action of Γ_3 to a point of the form

$$\begin{pmatrix} \tau_1 & 0 \\ 0 & \tau_2 \end{pmatrix}$$

with $\tau_i \in \mathcal{H}_{g_i}$, $g_1 + g_2 = 3$. We shall denote by $\mathcal{R}_3$ the set of reducible points, on which $\vartheta[0]$ vanishes. It is known that $\mathcal{R}_3/\Gamma_3[2]$ is contained in X and even more

$$X = X_0 \cup \overline{\mathcal{R}_3/\Gamma_3[2]} \quad (\text{disjoint union}).$$

We want to count how many reducible components appear in $X - X_0$. Each reducible component of $\overline{\mathcal{R}_3/\Gamma_3[2]}$ is defined by the vanishing of 6 thetanullwerte, (one of them zero, since we work on the locus $\vartheta[0] = 0$). We refer to [Gl] for details. A sextuplet of characteristics $m_1, \dots, m_6$ corresponding to these six thetanullwerte have the property that all sums of three of them result to be an odd characteristic. An example is given by

m_1	m_2	m_3	m_4	m_5	m_6
0	1	0	1	0	0
0	1	0	0	1	0
0	1	0	0	0	1
0	1	1	0	1	1
0	0	1	0	0	1
0	1	1	0	0	0

Hence we obtain:

7.1 Proposition. *The boundary part $X - X_0$ of the hyperelliptic component X consists of 56 irreducible 4-dimensional components. They correspond to sets of 6 even characteristics, one of them 0, such that all sums of three of them are odd. The component is the intersection of the corresponding zero divisors $\vartheta[m] = 0$. The group $\Gamma_{3,\vartheta}$ permutes the components transitively.*

Part of the boundary $X - X_0$ is the intersection of X with the Satake boundary. We want to describe this part more closely. Recall that

$$\mathcal{H}_3^* - \mathcal{H}_3$$

is the disjoint union of the Γ_3 -orbits of the three standard components

$$\begin{pmatrix} i\infty & 0 & 0 \\ 0 & i\infty & 0 \\ 0 & 0 & i\infty \end{pmatrix}, \quad \begin{pmatrix} \tau & 0 & 0 \\ 0 & i\infty & 0 \\ 0 & 0 & i\infty \end{pmatrix} \quad (\tau \in \mathcal{H}_1), \quad \begin{pmatrix} \tau & 0 \\ 0 & i\infty \end{pmatrix} \quad (\tau \in \mathcal{H}_2).$$

We denote by $\mathcal{D}_3$ the part of $\mathcal{H}_3^* - \mathcal{H}_3$ on which $\vartheta[0]$ vanishes. We see that $\mathcal{D}_3/\Gamma_3[2]$ is contained already in the closure of the reducible locus. Hence we have the inclusions

$$\mathcal{D}_3/\Gamma_3[2] \subset \overline{\mathcal{R}_3/\Gamma_3[2]} \subset X \subset \overline{\mathcal{H}_3/\Gamma_3[2]}.$$

Next we want to describe the zero dimensional boundary components of X . These are the points which come from the orbit of the point

$$\begin{pmatrix} i\infty & 0 & 0 \\ 0 & i\infty & 0 \\ 0 & 0 & i\infty \end{pmatrix}.$$

In [Gl] one finds the description of the zero dimensional boundary components as intersections of 28 zero divisors $\vartheta[m] = 0$. This description can be related to stars. Recall that a star is a set of 4 odd characteristics with a certain property (2.2).

7.2 Proposition. *Let S be a star. There are 27 even characteristics $m \neq 0$ orthogonal to at least one characteristic in S . The intersection of the 27 divisors $\vartheta[m] = 0$ is a zero-dimensional boundary component of X . This gives a 1-1-correspondence between the 105 stars and the 105 zero-dimensional boundary components.*

We mention also how the remaining 8 even characteristics can be described in a nice way: The star S spans a three dimensional space M . There exists a unique even characteristic n such that $M + n$ contains only even characteristics. The entries of this coset are the 8 remaining characteristics.

Now we describe the closures of the three dimensional boundary components of X . They are of the type

$$\begin{pmatrix} \tau & 0 \\ 0 & i\infty \end{pmatrix}, \quad \tau \in \mathcal{H}_2.$$

In [Gl] one finds also the description of the three dimensional boundary components as intersections of 16 zero divisors $\vartheta[m] = 0$. The description can be related to the odd characteristics

7.3 Proposition. *Let α be an odd characteristic, there are exactly 15 even characteristics $m \neq 0$ orthogonal to it. The corresponding divisors $\vartheta[m] = 0$ intersect in a three-dimensional boundary component of X . This gives a 1-1-correspondence between the 28 odd characteristics and the 28 irreducible components of the boundary $\mathcal{D}_3/\Gamma_3[2]$ which are all three dimensional.*

A similar computation can be done for the one dimensional boundary components of $\overline{\mathcal{H}_3/\Gamma_3[2]}$, knowing that in this case each component is given by the vanishing of 24 suitable thetanullwerte, cf [Gl]. In this case we get:

7.4 Proposition. *Let α_1 and α_2 be odd characteristics such that $\alpha_1 + \alpha_2$ is an even characteristics, then there are exactly 23 even characteristics $m \neq 0$ orthogonal to α_1 or to α_2 . The corresponding divisors $\vartheta[m] = 0$ intersect in a one-dimensional boundary component of X . This gives a 1-1-correspondence between the 210 pairs of odd characteristics of the above form and the 210 one components of the boundary $\mathcal{D}_3/\Gamma_3[2]$.*

Next we recall the structure of $Y - Y_0$. We have two possibilities, to use the GIT-picture or the ball-picture. The GIT-picture has been explained in [Koi]. We just recall the following: Consider the map

$$\mathbb{C}^8 \longrightarrow (P^1)^8 \longrightarrow (P^1)^8 // \mathrm{SL}(2) = Y.$$

In section one we used the variables $X_1, \dots, X_8$ to describe this $\mathbb{C}^8$. We have to consider the 28 differences $W_{ij} = X_i - X_j$, $1 \leq i < j \leq 8$. The images of their zero set in Y describe 28 irreducible divisors. Their union is the boundary part $Y - Y_0$. We now switch to the ball picture. The reason is that in this picture the quadratic space $\mathbb{F}_2^6$ occurs, whose elements also play the role of theta characteristics.

In 3.3 we showed that the 28 pairs (i, j) are in 1-1-correspondence to the anisotropic elements of $\mathbb{F}_2^6$, in other words to the odd characteristics. Recall that to each such element α we introduced an irreducible divisor (Heegner divisor) $\mathcal{B}_\alpha$ in $\overline{\mathcal{B}}/\Gamma[1 + i]$.

7.5 Proposition. *In our identification $Y = \overline{\mathcal{B}}/\Gamma[1 + i]$ the boundary divisor in Y defined by “ $W_{ij} = 0$ ”, cf. Proposition 3.2, corresponds to the Heegner divisor $\mathcal{B}_\alpha$, where α is the anisotropic element of $\mathbb{F}_2^6$ related to (i, j) as in 3.3.*

For a sequence of odd characteristics $\alpha_1, \dots, \alpha_k$, we associate the intersection

$$\mathcal{B}_{\alpha_1, \dots, \alpha_k} := \mathcal{B}_{\alpha_1} \cap \dots \cap \mathcal{B}_{\alpha_k}.$$

Now we want to compare the Siegel- and the ball picture, i.e. we have to consider the rational map $Y \dashrightarrow X$.

7.6 Proposition. *Under the rational map $Y \dashrightarrow X$ the 28 Heegner divisors are contracted to the 28 three-dimensional boundary components. The 210 intersections $\mathcal{B}_{\alpha_1, \alpha_2}$ with $\alpha_1 + \alpha_2$ an even characteristic are contracted to the 210 one dimensional boundary components. The 420 intersections $\mathcal{B}_{\alpha_1, \alpha_2, \alpha_3}$ with $\alpha_i + \alpha_j$ even characteristics ($1 \leq i < j \leq 3$) are contracted to the 105 zero dimensional boundary components. More precisely, the four $\mathcal{B}_{\alpha_1, \alpha_2, \alpha_3}$ obtained by a star map to the same point.*

The rational map is defined outside the base locus, which has been described in 6.2 as a concrete three dimensional variety. When B is a divisor in X we can take its inverse image in the complement of the base locus and then take the closure A in X . We call A the pull-back of B . From 3.2 and 3.3 we get:

7.7 Proposition. *The pull-back of the zero divisor of the $\vartheta[m]$, $m \neq 0$, is the union of 12 Heegner divisors $\mathcal{B}_\alpha$, namely those such that α is orthogonal to m .*

Recall (2.2) that stars consist of 4 anisotropic vectors with a certain property. The union of the four $\mathcal{B}_\alpha$ is the star-divisor $\mathcal{B}_S$. A direct inspection shows (one m is enough):

7.8 Remark. *Let $m \neq 0$ be an even characteristic. There are five different ways to write the set of all anisotropic α , which are orthogonal to m , as union of three stars. The pull-back of the zero divisor of $\vartheta[m]$ is the union of any of such three star divisors.*

We come now to a description of the cusps. It is known that there are 35 cusps (elements of $\overline{\mathcal{B}}/\Gamma[1-i] - \mathcal{B}/\Gamma[1-i]$). In the GIT-model they correspond to the instable points. From [Koi] or as an almost direct consequence of the computational Lemma 6.3, we obtain:

7.9 Remark. *The intersection of the twelve Heegner divisors, which correspond to an even non-zero characteristic (7.7), is a cusp. This gives a bijection between the cusps and the non-zero even characteristics.*

Now we will give a combinatorial description of the base locus of the rational map $Y \dashrightarrow X$. We recall that it is the union of 56 linear spaces. This has been proved in 6.2 by a computer calculation (in the strong ideal theoretic sense). Here will give an intrinsic (set-theoretical) description of this locus in terms of Heegner divisors.

First of all we need some preliminary facts.

7.10 Lemma. *Let α_1 and α_2 be two odd characteristics such that*

$$\alpha_3 = \alpha_1 + \alpha_2$$

is still an odd characteristic. Then each even characteristic $m \neq 0$ is orthogonal to at least one of the three $\alpha_1, \alpha_2, \alpha_3$.

In fact each odd characteristics α_1 is orthogonal to 15 even characteristic $m \neq 0$. Moreover 25 even characteristics $m \neq 0$ are orthogonal to at least one of two odd characteristics α_1 and α_2 whose sum is still odd, thus both are orthogonal to 5 even characteristics $m \neq 0$. Obviously $\alpha_3 = \alpha_1 + \alpha_2$ is still orthogonal to the same 5 even characteristics m , and there are 10 more that are orthogonal to α_3 , but neither orthogonal to α_1 nor to α_2 . So we get that each even characteristic m is orthogonal to at least one among α_1, α_2 and α_3 .

This discussion also shows:

7.11 Corollary. *There are exactly 5 even characteristics $m \neq 0$ orthogonal to $\alpha_1, \alpha_2, \alpha_3$ in 7.10.*

Let us consider a different configuration. We assume that α_1, α_2 are odd but $\alpha_1 + \alpha_2$ is an even characteristic. In this case 23 even non zero characteristics are orthogonal to at least one of α_1, α_2 , and both together are orthogonal to 7 even characteristics. Let us take an odd α_3 such that $\alpha_1 + \alpha_3$ and $\alpha_2 + \alpha_3$ are even characteristics. Then 27 non zero even characteristics are orthogonal to at least one among $\alpha_1, \alpha_2, \alpha_3$. There is one and only one odd characteristic α_4 which added to $\alpha_1, \alpha_2, \alpha_3$ gives even characteristics, namely $\alpha_4 = \alpha_1 + \alpha_2 + \alpha_3$. Hence the four odd characteristics $\alpha_1, \alpha_2, \alpha_3, \alpha_4$ form a star. From this discussion follows:

7.12 Lemma. *Let $\alpha_1, \dots, \alpha_k$ be a sequence of odd characteristics such that each even characteristic is orthogonal at least to a α_i . Then the sequence contains three characteristics, say $\alpha_1, \alpha_2, \alpha_3$, with $\alpha_3 = \alpha_1 + \alpha_2$.*

For three odd characteristics $\alpha_1, \alpha_2, \alpha_3$ with the property $\alpha_1 + \alpha_2 + \alpha_3 = 0$ we want to consider the intersection $\mathcal{B}_{\alpha_1, \alpha_2, \alpha_3}$. One can check that it coincides already with the intersection of any two of the three Heegner divisors, i.e

$$\mathcal{B}_{\alpha_1, \alpha_2, \alpha_3} = \mathcal{B}_{\alpha_1, \alpha_2} = \mathcal{B}_{\alpha_1, \alpha_3} = \mathcal{B}_{\alpha_2, \alpha_3}.$$

A direct computation tells us that there are exactly 56 triplets of the form $\alpha_1, \alpha_2, \alpha_3$, with $\alpha_1 + \alpha_2 + \alpha_3 = 0$. With these notations we have the following

7.13 Proposition. *The base locus of the rational map from Y to X is the set*

$$V = \bigcup_{\alpha_1 + \alpha_2 + \alpha_3 = 0} \mathcal{B}_{\alpha_1, \alpha_2, \alpha_3}.$$

To each $\mathcal{B}_{\alpha_1, \alpha_2, \alpha_3}$ corresponds to a linear space of Proposition 6.2.

The proof is an immediate consequence of the previous lemmata and of the fact that the base locus is necessarily the union of intersections of Heegner divisors.

We know from the results of the previous sections that the blow up of the base locus produces 56 divisors in Y^* or in $\bar{M}_{0,8}$. We are interested in the image of these divisors in X . From the previous corollary 7.11, we have exactly 5 even characteristics orthogonal to $\alpha_1, \alpha_2, \alpha_3$. Thus the image of $\mathcal{B}_{\alpha_1, \alpha_2, \alpha_3}$ is defined by the vanishing of 6 thetanullwerte whose characteristics are $\Gamma_{3, \vartheta}$ -conjugate to those at the begin of the section.

Thus, according to our previous result, we have that they define an irreducible component of $\overline{\mathcal{R}_3}/\Gamma_3[2]$. Hence we get:

7.14 Proposition. *The blow up of the base locus maps on the reducible locus of the hyperelliptic modular variety. In particular the blow up of $\mathcal{B}_{\alpha_1, \alpha_2, \alpha_3}$ maps onto the component defined by the vanishing of the thetanullwerte $\vartheta[m]$ with m orthogonal to $\alpha_1, \alpha_2, \alpha_3$.*

Let us conclude considering the image of the blow up of the cusps. First of all we observe that for each even, non zero characteristic m there are 16 even characteristics $n_1, \dots, n_{16}$ such that for each index $i = 1, \dots, 16$, $m + n_i = \beta_i$ is an odd characteristic. These sixteen odd characteristics appear in the situation that we are going to describe.

We recall from Lemma 6.3 and Proposition 7.13 that 8 linear components $\mathcal{B}_{\alpha_1, \alpha_2, \alpha_3}$ intersect in a cusp m . Each triplet is characterized by being orthogonal to the characteristic m . Viceversa, according to Corollary 7.11, to each triplet $\alpha_1, \alpha_2, \alpha_3$ we can associate, besides the characteristics m , four more even characteristics that are orthogonal to them. The union of all these even characteristics appearing at least once for the eight possible triplets $\alpha_1, \alpha_2, \alpha_3$, related to the even characteristic m , gives the sixteen even characteristics

$$n_1, \dots, n_{16}.$$

In [Gl] one finds also the description of the two dimensional components conjugate to

$$\begin{pmatrix} \tau_1 & 0 & 0 \\ 0 & \tau_2 & 0 \\ 0 & 0 & i\infty \end{pmatrix} \quad (\tau_i \in \mathcal{H}_1)$$

as intersections of 18 zero divisors $\vartheta[n] = 0$. The description can be related to the characteristics $0, m, n_1, \dots, n_{16}$. In fact we have

7.15 Proposition. *The blow up of the cusps maps on two dimensional components in the hyperelliptic modular variety. In particular the blow up of the cusp m maps onto the component defined by the vanishing of the thetanullwerte $\vartheta[n]$, $n = 0, m, n_1, \dots, n_{16}$ with $m + n_i$ odd.*

We remind that in Proposition 6.4 is named the two dimensional variety that is the inverse image of the cusps in Y^* .

References

- [AF] Allcock, D. and Freitag, E.: *Cubic surfaces and Borcherds products*, Commentarii Math. Helv. Vol. **77**, Issue 2, 270–296 (2002)
- [AL] Avritzer, D. Lange, H.: *The moduli spaces of hyperelliptic curves and binary forms*, Math. Z. **242**, 615–632 (2002)
- [Bo] Borcherds, R.: *Automorphic forms with singularities on Grassmannians*, Invent. math. **132**, 491–562 (1998)
- [Fr1] Freitag, E.: *Some modular forms related to cubic surfaces* Kyungpook Math. J. **43**, No.3, 433–462 (2003)
- [Fr2] Freitag, E.: *Comparison of different models of the moduli space of marked cubic surfaces*, Proceedings of Japanese-German Seminar, Ryushi-do, edited by T. Ibukiyama and W. Kohnen, 74–79 (2002)

- [Fr3] Freitag, E.: *Siegelsche Modulfunktionen* Grundlehren der mathematischen Wissenschaften, **254** Berlin-Heidelberg-New York: Springer-Verlag (1983)
- [FS] Freitag, E. Salvati-Manni, R.: *Modular forms for the even unimodular lattice of signature (2,10)*, J. Algebraic Geom. **16**, 753–791 (2007)
- [Gl] Glass, J. : *Theta constants of genus three*, Compos. Math. **40**, 123-137 (1980).
- [Ho] Howe, R.: *The classical groups and invariants of bilinear forms*, The Mathematical Heritage of Hermann Weyl (Durham, NC, 1987), 133–166, Proc. Sympos. Pure Math. **48**, Amer. Math. Soc., Providence, RI (1988)
- [HM1] Howard, B.J. Millson, J. Snowden, A. Vakil, R.: *The projective invariants of ordered points on the line*, ArXiv Mathematics e-prints math.AG/0505096 (2007)
- [HM2] Howard, B.J. Millson, J. Snowden, A. Vakil, R.: *The moduli space of n points on the line is cut out by simple quadrics when n is not six*, ArXiv Mathematics e-prints math.AG/0607372 (2007)
- [Ig1] Igusa, J.: *On the graded ring of theta-constants* Am. J. Math. **86**, 219-246 (1964).
- [Ig2] Igusa, J.: *Modular forms and projective invariants* Am. J. Math. **89**, 817-855 (1967).
- [Ka] Kapranov, M.M.: *Chow quotients of Grassmannians I*, Adv. Sov. Math. **16** (2), 29-110 (1993)
- [Ko1] Kondo, S.: *The moduli space of Enriques surfaces and Borcherds products*, J. Algebraic Geometry **11**, 601-627(2002)
- [Ko2] Kondo, S.: *The moduli space of 8 points on $P^1(\mathbb{C})$ and automorphic forms* in Algebraic Geometry Contemporary Mathematics **422**, 89-106 (2007), Amer. Math.Soc.
- [Koi] Koike, K.: *The projective embedding of the configuration space $X(2, 8)$* , Technical Reports of Mathematical Sciences, Chiba University, **16** (2000)
- [MY] Matsumoto, K. Yoshida, M.: *Configuration space of 8 points on the projective line and a 5-dimensional Picard modular group*, Compositio Math. **86**, 265–280 (1993)
- [Mu] *Tata Lectures on Theta II*, Modern Birkhuser Classics, Reprint of the 1984 ed., XIV (2007)
- [Ru1] Runge, B.: *On Siegel modular form, part I*, J. Reine angew. Math. **436**, 57-85 (1993)
- [Ru2] Runge, B.: *On Siegel modular forms, part II*, Nagoya Math. J. **138**, 179-197 (1995)
- [Ts] Tsuyumune, S.: *Thetanullwerte on a moduli space of curves and hyperelliptic loci*, Math. Zeit. **207**, 539-568 (1991)